

```
import pandas as pd
import numpy as np
import matplotlib.pyplot as plt
```

```
df = pd.read_csv("data.csv")
df.head()
```

	id	diagnosis	radius_mean	texture_mean	perimeter_mean	area_mean
0	842302	M	17.99	10.38	122.80	1001.0
1	842517	M	20.57	17.77	132.90	1326.0
2	84300903	M	19.69	21.25	130.00	1203.0
3	84348301	M	11.42	20.38	77.58	386.1
4	84358402	M	20.29	14.34	135.10	1297.0

5 rows × 33 columns

```
df.isnull().sum()
```


	0
id	0
<pre>df = df.drop(columns=["id", "Unnamed: 32"]) df.fillna(df.mean(numeric_only=True), inplace=True) df.isnull().sum()</pre>	
texture_mean	0
perimeter_mean	0
area_mean	0
smoothness_mean	0
compactness_mean	0
concavity_mean	0
concave points_mean	0
symmetry_mean	0
fractal_dimension_mean	0
radius_se	0
texture_se	0
perimeter_se	0
area_se	0
smoothness_se	0
compactness_se	0
concavity_se	0
concave points_se	0
symmetry_se	0
fractal_dimension_se	0
radius_worst	0
texture_worst	0
perimeter_worst	0
area_worst	0
smoothness_worst	0
compactness_worst	0
concavity_worst	0
concave points_worst	0
symmetry_worst	0
fractal_dimension_worst	0

dtype: int64


```
def remove_outliers(df, column):
    Q1 = df[column].quantile(0.25)
    Q3 = df[column].quantile(0.75)
    IQR = Q3 - Q1
    lower = Q1 - 1.5 * IQR
    upper = Q3 + 1.5 * IQR
    return df[(df[column] >= lower) & (df[column] <= upper)]
```

```
for col in df.select_dtypes(include=np.number).columns:
    df = remove_outliers(df, col)
```

```
df.shape
```

```
concavity_mean      0
(277, 31)
concave points_mean  0
```

```
plt.figure()
plt.hist(df['radius_mean'])
plt.show()
```



```
plt.figure()
plt.hist(df['area_mean'])
plt.show()
```