

Probability and Statistics

Prepared by Dr. Alfaisal Hasan

STA 201
Probability and Statistics

Lecture NO 4





مجموعة جامعة
الجميع الذكية
للتعليم القابضة

MIDOCEAN
UNIVERSITY

Probability and Statistics

Lecture NO 4

STA 201

Probability and Statistics

Prepared by
Dr. Alfaisal Hasan

www.midocean.ae

Chapter

Topics

- Chapter 1: Introduction To Statistics
- Chapter 2: Descriptive Statistics
- Chapter 3: Probability
- **Chapter 4: Discrete Probability Distributions**
- Chapter 5: Normal Probability Distributions
- Chapter 6: Confidence Intervals
- Chapter 7: Hypothesis Testing with One Sample
- Chapter 8: Hypothesis Testing with Two Samples
- Chapter 9: Correlation And Regression
- Chapter 10: Chi-Square Tests and The F -Distribution

Chapter

Discrete Probability Distributions

4.1 Probability Distributions

4.2 Binomial Distributions

4.3 More Discrete Probability Distributions

Discrete Probability Distributions

- Distinguish between discrete random variables and continuous random variables
- Construct a discrete probability distribution and its graph
- Determine if a distribution is a probability distribution
- Find the mean, variance, and standard deviation of a discrete probability distribution
- Find the expected value of a discrete probability distribution

Probability Distributions

Random Variable

Represents a numerical value associated with each outcome of a probability distribution.

Denoted by x

Examples

x = Number of sales calls a salesperson makes in one day.

x = Hours spent on sales calls in one day.


Discrete Random Variable

Discrete Random Variable

Has a finite or countable number of possible outcomes that can be listed.

Example:

x = Number of sales calls a salesperson makes in one day.




Continuous Random Variable

Continuous Random Variable

Has an uncountable number of possible outcomes, represented by an interval on the number line.

Example:

x = Hours spent on sales calls in one day.




Example: Random variable

Decide whether the random variable x is discrete or continuous.

x = The number of stocks in the Dow Jones Industrial Average that have share price increases on a given day.

Solution:

Discrete random variable (The number of stocks whose share price increases can be counted.)




Example: Random variable

Decide whether the random variable x is discrete or continuous.
 x = The volume of water in a 32-ounce container.

Solution:

Continuous random variable (The amount of water can be any volume between 0 ounces and 32 ounces)



Discrete Probability Distribution

Discrete probability distribution

Lists each possible value the random variable can assume, together with its probability.

Must satisfy the following conditions:

In Words

1. The probability of each value of the discrete random variable is between 0 and 1, inclusive.
2. The sum of all the probabilities is

In Symbols

$$0 \leq P(x) \leq 1$$

$$\Sigma P(x) = 1$$

Constructing a Discrete Probability Distribution

Let x be a discrete random variable with possible outcomes $x_1, x_2, \dots, x_n$.

1. Make a frequency distribution for the possible outcomes.
2. Find the sum of the frequencies.
3. Find the probability of each possible outcome by dividing its frequency by the sum of the frequencies.
4. Check that each probability is between 0 and 1 and that the sum is 1.

Example: Constructing a Discrete Probability Distribution

An industrial psychologist administered a personality inventory test for passive-aggressive traits to 150 employees. Individuals were given a score from 1 to 5, where 1 was extremely passive and 5 extremely aggressive. A score of 3 indicated neither trait. Construct a probability distribution for the random variable x .

Score, x	Frequency, f
1	24
2	33
3	42
4	30
5	21

Solution: Constructing a Discrete Probability Distribution

Divide the frequency of each score by the total number of individuals in the study to find the probability for each value of the random variable.

$$P(1) = \frac{24}{150} = 0.16 \quad P(2) = \frac{33}{150} = 0.22 \quad P(3) = \frac{42}{150} = 0.28$$

$$P(4) = \frac{30}{150} = 0.20 \quad P(5) = \frac{21}{150} = 0.14$$

- Discrete probability distribution:

x	1	2	3	4	5
$P(x)$	0.16	0.22	0.28	0.20	0.14

Solution: Constructing a Discrete Probability Distribution

x	1	2	3	4	5
$P(x)$	0.16	0.22	0.28	0.20	0.14




This is a valid discrete probability distribution since

1. Each probability is between 0 and 1, inclusive,
 $0 \leq P(x) \leq 1$
2. The sum of the probabilities equals 1,
 $\Sigma P(x) = 0.16 + 0.22 + 0.28 + 0.20 + 0.14 = 1$

Solution: Constructing a Discrete Probability Distribution

Histogram



Because the width of each bar is one, the area of each bar is equal to the probability of a particular outcome.

Mean, Variance and standard deviation for probability Distribution

Mean of a discrete probability distribution $\mu = \sum xP(x)$

Each value of x is multiplied by its corresponding probability and the products are added. Also, mean is known as (Expected Value).

Variance of a discrete probability distribution $\sigma^2 = \sum (x - \mu)^2 P(x)$

Standard deviation of a discrete probability distribution

$$\sigma = \sqrt{\sigma^2} = \sqrt{\sum (x - \mu)^2 P(x)}$$

Example:

The probability distribution for the personality inventory test for passive-aggressive traits is given.

Find the mean (Expected value)

Find Variance and Standard deviation

Solution:

$$\mu = \sum x P(x) = 2.94$$

x	$P(x)$	$xP(x)$
1	0.16	$1(0.16) = 0.16$
2	0.22	$2(0.22) = 0.44$
3	0.28	$3(0.28) = 0.84$
4	0.20	$4(0.20) = 0.80$
5	0.14	$5(0.14) = 0.70$

Mean, Variance and standard deviation

x	$P(x)$	$x - \mu$	$(x - \mu)^2$	$(x - \mu)^2 P(x)$
1	0.16	$1 - 2.94 = -1.94$	$(-1.94)^2 = 3.764$	$3.764(0.16) = 0.602$
2	0.22	$2 - 2.94 = -0.94$	$(-0.94)^2 = 0.884$	$0.884(0.22) = 0.194$
3	0.28	$3 - 2.94 = 0.06$	$(0.06)^2 = 0.004$	$0.004(0.28) = 0.001$
4	0.20	$4 - 2.94 = 1.06$	$(1.06)^2 = 1.124$	$1.124(0.20) = 0.225$
5	0.14	$5 - 2.94 = 2.06$	$(2.06)^2 = 4.244$	$4.244(0.14) = 0.594$

Variance

$$\sigma^2 = \Sigma(x - \mu)^2 P(x) = 1.616$$

standard deviation

$$\sigma = \sqrt{\sigma^2} = \sqrt{1.616} \approx 1.3$$

Section 4.2 Objectives

- Determine if a probability experiment is a binomial experiment
- Find binomial probabilities using the binomial probability formula
- Find binomial probabilities using technology and a binomial table
- Graph a binomial distribution
- Find the mean, variance, and standard deviation of a binomial probability distribution

Binomial Experiments

1. The experiment is repeated for a fixed number of trials, where each trial is independent of other trials.
2. There are only two possible outcomes of interest for each trial. The outcomes can be classified as a success (S) or as a failure (F).
3. The probability of a success $P(S)$ is the same for each trial.
4. The random variable x counts the number of successful trials.

Notation for Binomial Experiments

<i>Symbol</i>	<i>Description</i>
n	The number of times a trial is repeated
$p = P(s)$	The probability of success in a single trial
$q = P(F)$	The probability of failure in a single trial ($q = 1 - p$)
x	The random variable represents a count of the number of successes in n trials: $x = 0, 1, 2, 3, \dots, n$.

Binomial Probability Formula

Binomial Probability Formula

The probability of exactly x successes in n trials is

$$P(x) = C_x^n p^x q^{n-x} = \frac{n!}{(n-x)!x!} p^x q^{n-x}$$

- n = number of trials
- p = probability of success
- $q = 1 - p$ probability of failure
- x = number of successes in n trials

Example

Micro fracture knee surgery has a 75% chance of success on patients with degenerative knees. The surgery is performed on three patients. Find the probability of the surgery being successful on exactly two patients.

Solution:

$$n=3 \quad p= 75\% = 0.75 \quad q = 1- 0.75 = 0.25$$




$$C_x^n p^x q^{n-x} = \frac{n!}{(n-x)!x!} p^x q^{n-x}$$

Solution

$$\begin{aligned}
 P(x=2) &= C_x^n p^x q^{n-x} \\
 &= C_2^3 (0.75)^2 (0.25)^{3-2} \\
 &= \frac{3!}{(3-2)! 2!} (0.75)^2 (0.25)^{3-2} = 3 (0.75)^2 (0.25)^{3-2} = 0.422
 \end{aligned}$$

Example

In a survey, workers in the U.S. were asked to name their expected sources of retirement income. Seven workers who participated in the survey are randomly selected and asked whether they expect to rely on Social Security for retirement income and found it equal to 25%. Create a binomial probability distribution for the all values.



Solution

25% of working Americans expect to rely on Social Security for retirement income.

$$n = 7, \quad p = 0.25, \quad q = 0.75, \quad x = 0, 1, 2, 3, 4, 5, 6, 7$$

$$P(x=0) = {}_7C_0(0.25)^0(0.75)^7 = 1(0.25)^0(0.75)^7 \approx 0.1335$$

$$P(x=1) = {}_7C_1(0.25)^1(0.75)^6 = 7(0.25)^1(0.75)^6 \approx 0.3115$$

$$P(x=2) = {}_7C_2(0.25)^2(0.75)^5 = 21(0.25)^2(0.75)^5 \approx 0.3115$$

$$P(x=3) = {}_7C_3(0.25)^3(0.75)^4 = 35(0.25)^3(0.75)^4 \approx 0.1730$$

$$P(x=4) = {}_7C_4(0.25)^4(0.75)^3 = 35(0.25)^4(0.75)^3 \approx 0.0577$$

$$P(x=5) = {}_7C_5(0.25)^5(0.75)^2 = 21(0.25)^5(0.75)^2 \approx 0.0115$$

$$P(x=6) = {}_7C_6(0.25)^6(0.75)^1 = 7(0.25)^6(0.75)^1 \approx 0.0013$$

$$P(x=7) = {}_7C_7(0.25)^7(0.75)^0 = 1(0.25)^7(0.75)^0 \approx 0.0001$$

Example

A survey indicates that 41% of women in the U.S. consider reading their favorite leisure-time activity. You randomly select four U.S. women and ask them if reading is their favorite leisure-time activity. Find the probability that at least two of them respond yes.



Solution:

- $n = 4$, $p = 0.41$, $q = 0.59$
- At least two means two or more.
- Find the sum of $P(2)$, $P(3)$, and $P(4)$.

Solution

$$P(X=2) = {}_4C_2(0.41)^2(0.59)^2 = 6(0.41)^2(0.59)^2 \approx 0.351094$$

$$P(X=3) = {}_4C_3(0.41)^3(0.59)^1 = 4(0.41)^3(0.59)^1 \approx 0.162654$$

$$P(X=4) = {}_4C_4(0.41)^4(0.59)^0 = 1(0.41)^4(0.59)^0 \approx 0.028258$$

$$\begin{aligned} P(X \geq 2) &= P(2) + P(3) + P(4) \\ &\approx 0.351094 + 0.162654 + 0.028258 \\ &\approx 0.542 \end{aligned}$$



Mean, Variance and standard deviation for Binomial Distribution

- **Mean:** $\mu = np$
- **Variance:** $\sigma^2 = npq$
- **Standard Deviation:** $\sigma = \sqrt{npq}$

Example

In Pittsburgh, Pennsylvania, about 56% of the days in a year are cloudy. Find the mean, variance, and standard deviation for the number of cloudy days during the month of June. (*Source: National Climatic Data Center*)

Solution: $n = 30$, $p = 0.56$, $q = 0.44$

Mean: $\mu = np = 30 \cdot 0.56 = 16.8$

Variance: $\sigma^2 = npq = 30 \cdot 0.56 \cdot 0.44 \approx 7.4$

Standard Deviation:



4.3 More Discrete Probability Distributions

- Find probabilities using the geometric distribution
- Find probabilities using the Poisson distribution

Geometric distribution

Geometric distribution

A discrete probability distribution.

Satisfies the following conditions:

- 1- A trial is repeated until a success occurs.
- 2- The repeated trials are independent of each other.
- 3- The probability of success p is constant for each trial.
- 4- The probability that the first success will occur on trial x is

$$P(x) = p(q)^{x-1},$$


where $q = 1 - p$.

Example

From experience, you know that the probability that you will make a sale on any given telephone call is 0.23. Find the probability that your first sale on any given day will occur on your fourth or fifth sales call.

Solution:

- $P(\text{sale on fourth or fifth call}) = P(4) + P(5)$
- Geometric with $p = 0.23$, $q = 0.77$, $x = 4, 5$



Solution

- $P(4) = 0.23(0.77)^{4-1} \approx 0.105003$
- $P(5) = 0.23(0.77)^{5-1} \approx 0.080852$

$$\begin{aligned} P(\text{sale on fourth or fifth call}) &= P(4) + P(5) \\ &\approx 0.105003 + 0.080852 \\ &\approx 0.186 \end{aligned}$$



Poisson distribution

A discrete probability distribution.

Satisfies the following conditions:

- The experiment consists of counting the number of times an event, x , occurs in a given interval.
- The interval can be an interval of time, area, or volume.
- The probability of the event occurring is the same for each interval.
- The number of occurrences in one interval is independent of the number of occurrences in other intervals.

Poisson distribution

Conditions continued:

The probability of the event occurring is the same for each interval.

The probability of exactly x occurrences in an interval is

$$P(x) = \frac{\mu^x e^{-\mu}}{x!}$$

where $e \approx 2.71818$ and μ is the mean number of occurrences

Example

The mean number of accidents per month at a certain intersection is 3. What is the probability that in any given month four accidents will occur at this intersection?



Solution

Solution:

- Poisson with $x = 4, \mu = 3$

$$P(4) = \frac{3^4(2.71828)^{-3}}{4!} \approx 0.168$$

Un Solved Examples

Chapter

If in Binomial distribution $n = 32$, $p = 0.7$, $q = 0.3$ then Find mean μ and
Variance σ^2 and Standard Deviation σ

Un Solved Examples

Chapter

If in Geometric distribution $p = 0.2$, $q = 0.8$ then Find $P(4)$

Un Solved Examples

Chapter

Number of rooms in the hospital is a discrete random variable

Chapter Summary

Mean of a discrete probability distribution = Expected value = $\mu = \sum x P(x)$

Variance of a discrete probability distribution $\sigma^2 = \sum (x - \mu)^2 P(x)$

Standard deviation of a discrete probability distribution

Chapter Summary

Chapter

Binomial Distributions :

$$P(X) = C_x^n p^x q^{n-x}$$

Where; $q = 1-p$



- **Mean:** $\mu = np$
- **Variance:** $\sigma^2 = npq$
- **Standard Deviation:** $\sigma = \sqrt{npq}$

Chapter Summary

Chapter

Geometric Distribution:

$$P(X) = p q^{x-1}$$

Poisson Distribution:

$$P(x) = \frac{\mu^x e^{-\mu}}{x!}$$



مجموعة جامعة
الجميع الذكية
للتعليم القابضة

MIDOCEAN
UNIVERSITY



Thank You

