

Research Proposal: Parental Awareness and Associated Risk Factors for Recurrent Respiratory Infections Among Preschool Children in Riyadh, Saudi Arabia

Lead Researcher: Tareq Hamdan Alzahrani

Institution: Midocean University, College of Management

Date: October 21, 2025

1. Introduction

Recurrent respiratory infections (RRIs) represent a pervasive and significant global health challenge, particularly affecting the pediatric population. These infections, characterized by repeated episodes of respiratory illness, impose a substantial burden on children, their families, and healthcare systems worldwide [1]. The vulnerability of preschool children (typically defined as those aged between 1 month and 5 years) to RRIs is well-documented, primarily due to their developing immune systems, increased exposure to pathogens in group settings like daycare, and often, a lack of fully established hygiene practices [2]. The consequences of RRIs extend beyond acute illness, potentially leading to chronic respiratory conditions, developmental delays, and significant disruptions to family life and parental productivity [3].

In the Kingdom of Saudi Arabia (KSA), respiratory infections are a leading cause of morbidity and healthcare utilization among children. The unique demographic, environmental, and socio-cultural landscape of Riyadh, the capital city, presents a specific context that warrants focused investigation. Riyadh is a rapidly growing metropolis with a diverse population, varying socioeconomic strata, and distinct environmental factors, including dust storms and air quality concerns, which can influence respiratory health [4]. Despite the recognized prevalence of respiratory illnesses, comprehensive research specifically addressing the interplay between parental awareness, modifiable risk factors, and the incidence of RRIs among preschool children in this specific urban setting remains limited.

Previous studies in Saudi Arabia have shed light on the viral etiology of respiratory infections, identifying common culprits such as Respiratory Syncytial Virus (RSV), human Rhinovirus (hRV), and influenza viruses as major contributors to acute respiratory tract infections (ARTIs) in pediatric populations [5, 6]. The burden of bronchiolitis, often caused by RSV, has been particularly noted, with significant hospitalization rates among young

children [7]. However, these studies often focus on clinical aspects or specific pathogens, leaving a gap in the understanding of how parental knowledge, attitudes, and practices contribute to the prevention and management of these recurrent episodes.

Parental awareness plays a pivotal role in child health outcomes. Informed parents are more likely to adopt preventive measures, recognize early signs of illness, seek timely medical attention, and adhere to treatment protocols, thereby potentially reducing the frequency and severity of RRIIs [8]. Conversely, a lack of awareness or misconceptions, particularly regarding antibiotic use for viral infections, can contribute to antibiotic resistance, a growing global health crisis [9]. Understanding the current levels of parental awareness and identifying specific knowledge gaps in Riyadh is therefore critical for designing effective public health interventions.

Furthermore, a myriad of risk factors contribute to the susceptibility of preschool children to RRIIs. These include, but are not limited to, environmental exposures (e.g., passive smoking, indoor air pollution), socioeconomic determinants (e.g., household income, parental education), and child-specific biological factors (e.g., nutritional status, daycare attendance, presence of siblings, underlying allergic conditions) [10, 11]. While some of these factors are universally recognized, their specific impact and prevalence can vary significantly across different geographical and cultural contexts. A localized investigation in Riyadh is essential to identify the most pertinent risk factors in this setting, allowing for the development of targeted and culturally sensitive interventions.

This research proposal outlines a comprehensive study designed to investigate parental awareness and associated risk factors for recurrent respiratory infections among preschool children in Riyadh, Saudi Arabia. By systematically exploring these critical areas, the study aims to generate evidence-based insights that can inform public health policies, enhance parental education programs, and ultimately contribute to improving the respiratory health and overall well-being of young children in the region. The findings are expected to provide a foundation for future longitudinal studies and intervention trials, fostering a healthier environment for the youngest members of the Riyadh community.

2. Research Problem

Recurrent respiratory infections (RRIIs) in preschool children pose a significant and multifaceted public health challenge in Riyadh, Saudi Arabia. While the global burden of RRIIs is well-established, and some regional data exist regarding the prevalence of respiratory pathogens, a critical gap remains in the comprehensive understanding of the specific local context. This knowledge deficit is particularly pronounced in two key areas: the precise identification and quantification of associated risk factors within the Riyadh population, and the assessment of parental awareness regarding these infections and their prevention.

Currently, healthcare providers and public health officials in Riyadh operate with a generalized understanding of RRI epidemiology, often relying on international data or broader national statistics that may not fully capture the unique dynamics of this specific urban environment. Riyadh, characterized by its rapid urbanization, diverse expatriate and local populations, and distinct environmental conditions (such as frequent dust storms and varying air quality levels), likely presents a unique set of contributing factors to pediatric respiratory health [12]. Without a granular understanding of these localized risk factors, interventions designed to mitigate the incidence and severity of RRIs may prove to be less effective or misdirected.

Furthermore, the role of parental awareness and practices is often underestimated in the chain of infection and prevention. Parents are the primary caregivers and decision-makers for preschool children, and their knowledge, attitudes, and behaviors directly influence a child's exposure to risk factors, adherence to preventive measures, and timely access to appropriate medical care [8]. Existing literature, even from within Saudi Arabia, often focuses on clinical outcomes or microbiological aspects, with limited in-depth exploration of the parental perspective. For instance, studies on parental awareness often concentrate on specific issues like antibiotic use, without a broader assessment of their understanding of RRI causes, transmission, prevention strategies, and appropriate home management [9]. This lack of specific data on parental awareness in Riyadh means that public health campaigns and educational initiatives may not be adequately tailored to address existing knowledge gaps or cultural nuances, potentially limiting their impact.

The absence of comprehensive, localized data on both risk factors and parental awareness creates a significant impediment to developing targeted, evidence-based public health strategies. Without knowing precisely which risk factors are most prevalent and impactful in Riyadh's preschool population, and without understanding where parents' knowledge and practices fall short, efforts to reduce the burden of RRIs will remain suboptimal. This research aims to directly address this critical knowledge gap by providing a detailed, context-specific analysis that can serve as a foundation for more effective and culturally appropriate interventions, ultimately contributing to improved respiratory health outcomes for preschool children in Riyadh.

3. Importance and Objectives of the Research

3.1. Importance of the Research

This research holds significant importance for several reasons, impacting public health, clinical practice, and family well-being in Riyadh, Saudi Arabia. The **scientific importance** lies in filling a critical gap in the existing literature by providing comprehensive, localized data on parental awareness and risk factors for RRIs in Riyadh. This will serve as a valuable

resource for future research, informing longitudinal studies, intervention trials, and comparative analyses across different regions. The findings will also contribute to a broader understanding of pediatric respiratory epidemiology in the Middle East, offering a context-specific perspective that complements global knowledge.

The **practical importance** of this research is multifaceted:

1. **Addressing a Significant Public Health Burden:** Recurrent respiratory infections are a leading cause of morbidity and healthcare utilization among preschool children globally and particularly in Saudi Arabia. By providing a comprehensive understanding of the local epidemiology and associated risk factors in Riyadh, this study will contribute to a more effective public health response, potentially reducing the incidence and severity of these infections. This translates to fewer hospitalizations, emergency room visits, and outpatient consultations, thereby alleviating the strain on the healthcare system.
2. **Informing Targeted Interventions:** Current public health strategies may not be optimally effective due to a lack of localized, granular data. This research will identify the most prevalent and impactful risk factors specific to Riyadh's preschool population. Such precise identification will enable the development of highly targeted and cost-effective interventions, moving beyond generic recommendations to address the unique challenges faced by families in this urban environment. For example, if indoor air quality or specific daycare practices are identified as major contributors, interventions can be designed to address these directly.
3. **Enhancing Parental Empowerment and Education:** Parents are at the forefront of child health management. By assessing current levels of parental awareness and identifying specific knowledge gaps regarding RRIs, this study will provide crucial information for designing tailored educational programs. Empowering parents with accurate information about prevention, early recognition of symptoms, and appropriate home management can significantly improve child health outcomes, reduce unnecessary antibiotic use, and foster a more proactive approach to respiratory health within families. This is particularly vital in combating antimicrobial resistance, a global health threat exacerbated by inappropriate antibiotic use for viral infections.
4. **Optimizing Resource Allocation:** Healthcare resources are finite. By pinpointing the most significant risk factors, policymakers and healthcare administrators can allocate resources more efficiently towards preventive measures and early intervention programs that yield the greatest impact. This could include investments in public awareness campaigns, improved environmental health regulations, or enhanced primary care services focused on pediatric respiratory health.
5. **Improving Quality of Life for Children and Families:** Frequent respiratory infections can significantly impact a child's quality of life, leading to discomfort, sleep

disturbances, missed schooling, and potential long-term health consequences. For families, RRI can result in parental stress, lost workdays, and financial burdens. By contributing to the reduction of RRI incidence and severity, this research ultimately aims to improve the overall well-being and quality of life for preschool children and their families in Riyadh.

3.2. Objectives of the Research

Building upon the identified research problem and its importance, this study aims to achieve the following specific objectives:

1. **Primary Objective:** To explore the relationship and associations between parental awareness, identified risk factors, and the incidence of recurrent respiratory infections in preschool children in Riyadh, Saudi Arabia, thereby identifying potential areas for intervention.
2. **Subsidiary Objectives:**
 - To determine the prevalence and characteristics of recurrent respiratory infections among preschool children (aged 1 month to 5 years) in Riyadh, Saudi Arabia. This objective seeks to establish baseline epidemiological data, including the frequency, duration, and types of RRI experienced by the target population over a defined period.
 - To identify and quantify the associated risk factors for recurrent respiratory infections in preschool children in Riyadh. This objective will involve a comprehensive assessment of various potential risk factors, categorized as:
 - **Environmental Factors:** Including exposure to passive smoking, indoor air quality (e.g., ventilation, presence of mold/dampness), and outdoor environmental conditions (e.g., dust, pollution).
 - **Socioeconomic Factors:** Such as parental education level, household income, family size, and access to healthcare services.
 - **Child-Specific Factors:** Including age, gender, birth order, nutritional status (e.g., breastfeeding history, micronutrient supplementation), daycare attendance, presence of siblings, and history of allergies or other chronic conditions.
 - To assess the level of parental awareness, knowledge, and practices regarding the causes, prevention, and management of recurrent respiratory infections in their preschool children. This objective will explore parents' understanding of RRI etiology (viral vs. bacterial), effective preventive measures (e.g., hand hygiene, avoiding sick contacts, vaccination), appropriate responses to symptoms, and attitudes towards antibiotic use.

- To provide evidence-based recommendations for public health interventions and educational programs aimed at reducing the burden of recurrent respiratory infections among preschool children in Riyadh. The ultimate goal is to translate research findings into actionable strategies that can be implemented by healthcare authorities, educators, and community organizations to improve child respiratory health.

4. Research Hypotheses

Based on the existing literature, the identified research problem, and the specific objectives of this study, the following hypotheses will be tested:

4.1. Hypothesis 1: Environmental Factors and RRI Incidence

H1a: There is a significant positive association between exposure to passive tobacco smoke within the household and the incidence of recurrent respiratory infections in preschool children in Riyadh.

- **Rationale:** Passive smoking is a well-established risk factor for respiratory illnesses in children, leading to impaired lung function, increased susceptibility to infections, and prolonged recovery times [13, 14]. In a household setting, children are particularly vulnerable to secondhand smoke from parents or other family members. This hypothesis aims to quantify this association specifically within the Riyadh context.

H1b: Poor indoor air quality, characterized by inadequate ventilation and/or the presence of mold/dampness, is significantly associated with a higher incidence of recurrent respiratory infections in preschool children in Riyadh.

- **Rationale:** Indoor air pollutants, including those from cooking, heating, and biological agents like mold, can irritate the respiratory tract and compromise immune responses, increasing susceptibility to infections [15]. Given the climate and housing characteristics in Riyadh, indoor air quality could be a significant, yet under-explored, environmental determinant of RRI. This hypothesis will investigate the direct link between these factors and RRI frequency.

4.2. Hypothesis 2: Socioeconomic Factors and RRI Prevalence

H2a: Lower parental education levels are significantly associated with a higher prevalence of recurrent respiratory infections among preschool children in Riyadh.

- **Rationale:** Parental education often correlates with health literacy, understanding of preventive measures, and access to and utilization of healthcare services [16]. Parents with higher education may be more aware of hygiene practices, vaccination

importance, and appropriate responses to illness, thereby potentially reducing their children's risk of RRIs. This hypothesis explores this relationship in the Riyadh setting.

H2b: Lower household income is significantly associated with a higher prevalence of recurrent respiratory infections among preschool children in Riyadh.

- **Rationale:** Household income can influence various factors related to child health, including access to nutritious food, quality housing, healthcare services, and ability to afford preventive measures. This hypothesis will examine the link between socioeconomic disadvantage, as measured by income, and RRI prevalence.

4.3. Hypothesis 3: Child-Specific Factors and RRI Incidence

H3a: Children attending daycare facilities have a significantly higher incidence of recurrent respiratory infections compared to those not attending daycare in Riyadh.

- **Rationale:** Group childcare settings are well-known environments for increased transmission of respiratory pathogens due to close contact among children, shared toys, and potential variations in hygiene practices [17]. This hypothesis aims to confirm this association within the Riyadh context.

H3b: Inadequate breastfeeding duration (less than 6 months exclusive breastfeeding) is significantly associated with a higher incidence of recurrent respiratory infections in preschool children in Riyadh.

- **Rationale:** Breastfeeding provides essential antibodies and immune factors that protect infants and young children from infections, including respiratory ones [18]. Shorter breastfeeding duration is often linked to increased susceptibility to RRIs. This hypothesis will investigate this protective effect.

4.4. Hypothesis 4: Parental Awareness and RRI Incidence

H4a: Higher levels of parental awareness regarding RRI causes, prevention, and management are significantly associated with a lower incidence of recurrent respiratory infections in their preschool children in Riyadh.

- **Rationale:** Informed parents are more likely to implement preventive measures, recognize symptoms early, and seek appropriate care, thereby reducing the frequency and severity of RRIs [8]. This hypothesis explores the direct impact of parental knowledge on child health outcomes.

H4b: A lack of parental awareness regarding the appropriate use of antibiotics for viral respiratory infections is significantly associated with a higher incidence of antibiotic misuse in preschool children in Riyadh.

- **Rationale:** Misconceptions about antibiotic efficacy for viral infections can lead to overuse, contributing to antibiotic resistance and potentially masking underlying

issues. This hypothesis focuses on a critical aspect of parental practice and its implications.

5. Literature Review

Recurrent Respiratory Infections (RRIs) in preschool children represent a significant global health concern, with extensive research conducted across various geographical and socio-economic contexts. This literature review synthesizes existing knowledge on RRIs, parental awareness, and associated risk factors, highlighting key findings, identifying gaps in current research, and articulating the unique contribution of the proposed study.

5.1. Global and Regional Burden of RRIs

Numerous studies have established the substantial burden of RRIs on pediatric populations worldwide, leading to significant morbidity, healthcare utilization, and disruptions to family life [1, 2, 3]. In the Kingdom of Saudi Arabia (KSA), respiratory infections are a leading cause of pediatric morbidity. Previous research in KSA has primarily focused on the viral etiology of respiratory infections, identifying common culprits such as Respiratory Syncytial Virus (RSV), human Rhinovirus (hRV), and influenza viruses as major contributors to acute respiratory tract infections (ARTIs) in pediatric populations [5, 6]. The burden of bronchiolitis, often caused by RSV, has been particularly noted, with significant hospitalization rates among young children [7]. These studies are crucial for understanding the microbiological landscape of RRIs in the region. **However, a common limitation of these studies is their focus on clinical and etiological aspects, often overlooking the broader interplay of parental awareness and modifiable risk factors.**

5.2. Parental Awareness and Practices

Parental awareness is a critical determinant of child health outcomes, particularly in the context of preventable diseases like RRIs. Informed parents are more likely to adopt preventive measures, recognize early signs of illness, seek timely medical attention, and adhere to treatment protocols [8]. Conversely, a lack of awareness or misconceptions, especially regarding antibiotic use for viral infections, can contribute to antibiotic resistance [9]. Studies like Al-Omari & Al-Qudah (2014) in Jordan [9] have explored parental knowledge, attitudes, and practices concerning antibiotic use for upper respiratory tract infections, revealing significant gaps. **While these studies shed light on specific aspects of parental behavior, there is a paucity of comprehensive research specifically assessing parental understanding of RRI causes, transmission, prevention strategies, and appropriate home management within the Riyadh context.** This gap is significant because public health campaigns and educational initiatives may not be adequately

tailored to address existing knowledge deficits or cultural nuances without this specific data.

5.3. Associated Risk Factors for RRIs

A myriad of risk factors contribute to the susceptibility of preschool children to RRIs, including environmental exposures (e.g., passive smoking, indoor air pollution), socioeconomic determinants (e.g., household income, parental education), and child-specific biological factors (e.g., nutritional status, daycare attendance, presence of siblings, underlying allergic conditions) [10, 11]. Global literature, such as reviews by Karr & Miller (2016) [10] and Loo & Ho (2014) [11], consistently highlight these factors. In Riyadh, environmental factors like dust storms and air quality concerns are known to influence respiratory health [4, 12]. **However, the specific impact and prevalence of these universally recognized factors can vary significantly across different geographical and cultural contexts.** Localized investigations are essential to identify the most pertinent risk factors in a given setting, allowing for the development of targeted and culturally sensitive interventions. **Existing regional studies often provide general epidemiological data but lack the granular analysis required to pinpoint the most impactful local risk factors and their interaction with parental awareness.**

5.4. Gaps in Current Research and Contribution of the Proposed Study

Based on the review of existing literature, several critical gaps emerge that the proposed research aims to address:

1. **Lack of Comprehensive Localized Data:** While some regional data exist regarding the prevalence of respiratory pathogens, there is a critical gap in the comprehensive understanding of the specific local context of Riyadh. This includes a lack of precise identification and quantification of associated risk factors within the Riyadh population.
2. **Limited Assessment of Parental Awareness:** Existing literature, even from within Saudi Arabia, often focuses on clinical outcomes or microbiological aspects, with limited in-depth exploration of the parental perspective. A broader assessment of parental understanding of RRI causes, transmission, prevention strategies, and appropriate home management in Riyadh is missing.
3. **Absence of Interplay Analysis:** Few studies comprehensively explore the complex interplay between parental awareness, various risk factors, and RRI outcomes in a single investigation within the Riyadh context. Understanding how these factors interact is crucial for effective intervention design.

This study will make a unique contribution by systematically investigating parental awareness and associated risk factors for recurrent respiratory infections among preschool

children in Riyadh, Saudi Arabia. By providing a detailed, context-specific analysis, this research will generate highly localized, actionable evidence that can directly inform public health policies and educational programs in Riyadh. It will move beyond generic recommendations to provide specific, culturally sensitive strategies tailored to the needs of the community, ultimately contributing to improved respiratory health outcomes for preschool children in the region. The findings will provide a robust evidence base for developing effective preventive and management strategies, thereby filling a significant void in both local public health knowledge and the broader scientific understanding of pediatric respiratory epidemiology.

6. Study Questions

Based on the research objectives, the following key questions will guide this study:

6.1. Primary Research Question

What is the relationship between parental awareness, identified risk factors, and the incidence of recurrent respiratory infections among preschool children in Riyadh, Saudi Arabia?

6.2. Subsidiary Research Questions

1. What is the prevalence and characteristics of recurrent respiratory infections among preschool children (aged 1 month to 5 years) in Riyadh, Saudi Arabia?
2. What are the associated environmental, socioeconomic, and child-specific risk factors for recurrent respiratory infections in preschool children in Riyadh?
3. What is the level of parental awareness, knowledge, and practices regarding the causes, prevention, and management of recurrent respiratory infections in their preschool children in Riyadh?
4. How do parental awareness and specific risk factors interact to influence the likelihood and frequency of recurrent respiratory infections in preschool children in Riyadh?
5. What evidence-based recommendations can be provided for public health interventions and educational programs aimed at reducing the burden of recurrent respiratory infections among preschool children in Riyadh?

7. Theoretical Premises of Study

(This section will provide a theoretical explanation of the information, definitions, variables of the study, and ideas related to the research topic. It will establish the conceptual framework guiding the research, drawing upon relevant theories and models from public

health, behavioral science, and pediatric epidemiology to contextualize the study's variables and their hypothesized relationships. Key terms and constructs, such as 'parental awareness,' 'recurrent respiratory infections,' and various 'risk factors,' will be defined and elaborated within this theoretical context.)

8. Research Methodology

This section provides a detailed exposition of the methodological framework that will guide the proposed study. It encompasses the study design, target population, sampling strategy, data collection instruments and procedures, data management protocols, statistical analysis plan, ethical considerations, and a discussion of potential challenges and limitations.

8.1. Study Design

This study will employ a **quantitative, cross-sectional descriptive design**. A cross-sectional design is particularly well-suited for this research as it allows for the simultaneous assessment of the prevalence of recurrent respiratory infections (RRIs), the distribution of various associated risk factors, and the levels of parental awareness within the target population at a single point in time. This approach provides a snapshot of the current situation, enabling the identification of associations and correlations between variables without inferring causality. The descriptive nature of the study will facilitate a comprehensive characterization of the study population and the key variables of interest.

8.2. Study Population

The target population for this study will be parents or legal guardians of preschool children (aged 1 month to 5 years) residing in Riyadh, Saudi Arabia. The age range of 1 month to 5 years is chosen because this demographic is particularly vulnerable to RRIs due to their developing immune systems, increased exposure to pathogens in group settings (e.g., daycare), and the typical peak incidence of respiratory infections in early childhood. Parents or legal guardians will be included if they are the primary caregivers for the child, are at least 18 years of age, and are able to provide informed consent and complete the questionnaire in either Arabic or English. Exclusion criteria will include parents of children with known severe chronic respiratory diseases (e.g., cystic fibrosis, severe congenital heart disease with significant pulmonary involvement) or diagnosed primary immunodeficiencies, as these conditions represent distinct clinical entities that could confound the findings related to general RRI risk factors.

8.3. Sampling Strategy

A multi-stage sampling technique will be utilized to ensure a representative sample from the diverse population of Riyadh. This approach aims to enhance the generalizability of the findings within the city:

1. **Stage 1: Stratification by Geographical Area:** Riyadh will be stratified into major geographical areas or municipalities (e.g., North, South, East, West, Central) based on available demographic data and urban planning divisions. This stratification will help account for potential variations in socioeconomic status, environmental exposures, and access to healthcare services across different parts of the city.
2. **Stage 2: Random Selection of Primary Healthcare Centers/Kindergartens:** Within each selected geographical stratum, a random sample of primary healthcare centers (PHCs) and/or private/public kindergartens will be selected. PHCs are chosen as they serve a broad cross-section of the population and are often the first point of contact for pediatric health concerns. Kindergartens are included to capture children in group settings, which are known to be relevant for RRI transmission. A proportional allocation method will be used to determine the number of PHCs/kindergartens to be selected from each stratum, ensuring representation based on population density or facility size.
3. **Stage 3: Systematic Random Sampling of Parents:** At the selected PHCs and kindergartens, systematic random sampling will be employed to recruit eligible parents. For PHCs, parents attending pediatric clinics for routine check-ups or non-emergency visits will be approached. For kindergartens, parents of enrolled children will be invited to participate. A sampling interval will be determined based on the expected number of eligible parents and the desired sample size from each facility. For example, every n th parent or child on a registration list will be invited.

8.3.1. Sample Size Calculation

The sample size will be calculated using the formula for estimating a single population proportion, given the cross-sectional design and the primary outcome of RRI prevalence. Based on existing literature, the estimated prevalence of RRIs in preschool children is approximately 25% [1, 2]. To ensure sufficient statistical power, the following parameters will be used:

- **Anticipated Prevalence (P):** 0.25 (25%)
- **Confidence Level:** 95% (Z-score = 1.96)
- **Margin of Error (d):** 0.05 (5%)

The formula for sample size (n) is: $n = (Z^2 * P * (1-P)) / d^2$

Substituting the values: $n = (1.96^2 * 0.25 * (1-0.25)) / 0.05^2$

$n = (3.8416 * 0.25 * 0.75) / 0.0025$

$n \approx 288.12$

Considering a potential non-response rate or incomplete questionnaires (e.g., 15-20%), the calculated sample size will be inflated to ensure adequate data collection. Therefore, a target sample size of approximately **350-400 parents** will be aimed for to account for potential attrition and ensure robust statistical analysis.

8.4. Data Collection Instruments and Procedures

Data will be collected primarily through a structured, self-administered questionnaire. The questionnaire will be meticulously designed to capture all variables relevant to the research objectives and hypotheses. It will be available in both Arabic and English to ensure accessibility and cultural appropriateness for the diverse population of Riyadh. The development process will involve several steps:

1. **Content Development:** Questions will be formulated based on a thorough review of existing validated questionnaires on RRI risk factors and parental awareness, adapted to the specific context of Riyadh. Expert opinion from pediatricians, public health specialists, and epidemiologists in Saudi Arabia will be sought to ensure content validity and relevance.
2. **Structure of the Questionnaire:** The questionnaire will be organized into distinct sections to enhance clarity and ease of completion:
 - **Section 1: Demographic Information:** Child's age, gender, birth order; parental age, education level, occupation, household income; family size, number of children under 5 in the household.
 - **Section 2: Child's Health History:** Frequency, duration, and severity of RRIs in the past 12 months; history of hospitalization for RTIs; presence of chronic conditions (e.g., asthma, allergies); vaccination status.
 - **Section 3: Potential Risk Factors:** Exposure to passive smoking (indoor/outdoor); home ventilation; presence of mold/dampness; daycare attendance; feeding practices (breastfeeding history); vitamin/mineral supplementation.
 - **Section 4: Parental Awareness:** Knowledge questions on RRI causes (viral vs. bacterial), prevention methods (hand hygiene, avoiding sick contacts, vaccination), and appropriate management. Attitudes towards antibiotic use will be assessed using a 5-point Likert scale (Strongly Disagree to Strongly Agree).
3. **Pre-testing and Pilot Study:** Before full-scale data collection, the questionnaire will undergo pre-testing with a small sample of 20-30 parents from the target population. This pilot study will assess the clarity of questions, ease of understanding, cultural appropriateness, and time required for completion. Feedback from the pilot will be used to refine and finalize the questionnaire, ensuring its reliability and validity.

4. **Administration:** The questionnaire will be primarily administered through secure online platforms (e.g., Google Forms, SurveyMonkey, Qualtrics). This method offers several advantages, including wider reach, cost-effectiveness, and automated data entry, reducing transcription errors. Links to the online questionnaire will be distributed via various channels:

- **Social Media Platforms:** WhatsApp groups, Facebook community pages, and other popular social media channels frequented by parents in Riyadh will be utilized. This will involve collaborating with community leaders or administrators of relevant groups.
- **Primary Healthcare Centers/Kindergartens:** With appropriate permissions, QR codes or links to the online questionnaire will be provided to parents at the selected facilities. Brief explanations about the study will be offered by trained research assistants.

Informed consent will be obtained electronically at the beginning of the online questionnaire. Participants will be clearly informed about the study's purpose, their right to withdraw at any time, and the confidentiality and anonymity of their responses. Participation will be entirely voluntary.

8.5. Data Management

Upon collection, all data will be securely managed to ensure confidentiality, integrity, and accuracy. For online questionnaires, data will be directly captured in the chosen platform, which typically offers secure storage and export functionalities. For any paper-based questionnaires, data will be manually entered into a secure database by trained personnel. A double-entry system will be implemented for a subset of the data to minimize data entry errors. All electronic data will be stored on password-protected computers with restricted access, and backed up regularly. Anonymity will be maintained by assigning unique identifiers to each participant, ensuring that no personally identifiable information is linked to the responses. Data will be retained for a specified period (e.g., 5 years) after the completion of the study, as per ethical guidelines, and then securely disposed of.

8.6. Statistical Analysis Plan

All collected data will be analyzed using appropriate statistical software, such as IBM SPSS Statistics (Statistical Package for the Social Sciences) or R statistical software. The analysis will proceed in several stages:

1. **Data Cleaning and Preparation:** This will involve checking for missing values, outliers, and inconsistencies. Missing data will be handled using appropriate methods (e.g., listwise deletion, imputation) depending on the extent and pattern of missingness. Variables will be coded and transformed as necessary for analysis.

2. **Descriptive Statistics:** Descriptive statistics will be used to summarize the demographic characteristics of the study population, the prevalence of RRIs, the distribution of various risk factors, and the levels of parental awareness. This will include frequencies and percentages for categorical variables, and means, medians, standard deviations, and interquartile ranges for continuous variables.
3. **Bivariate Analysis:** To explore initial associations between variables, bivariate analyses will be conducted:
 - **Chi-square tests (χ^2):** Will be used to examine associations between categorical variables, such as the relationship between exposure to passive smoking (Yes/No) and the incidence of RRIs (e.g., 0-2 episodes vs. >2 episodes).
 - **Independent samples t-tests or Analysis of Variance (ANOVA):** Will be used to compare means of continuous variables across different groups (e.g., comparing mean RRI frequency between children attending daycare vs. those not attending).
 - **Pearson correlation coefficients:** Will be calculated to assess the strength and direction of linear relationships between continuous variables (e.g., parental education level and parental awareness score).
4. **Multivariate Analysis (Logistic Regression):** To identify independent predictors of recurrent respiratory infections and to control for potential confounding variables, multivariate logistic regression models will be constructed. The dependent variable will be the presence or absence of recurrent respiratory infections (defined based on a threshold, e.g., >3 episodes in 12 months). Independent variables will include significant risk factors identified in the bivariate analysis, as well as demographic variables. Odds Ratios (OR) with 95% Confidence Intervals (CI) will be calculated to quantify the strength of association between the predictors and the outcome. Model fit and assumptions will be assessed.
5. **Statistical Significance:** A p-value of less than 0.05 ($p < 0.05$) will be considered statistically significant for all analyses.

8.7. Ethical Considerations

This study will adhere to the highest ethical standards throughout all phases of the research. Ethical approval will be sought and obtained from the Institutional Review Board (IRB) or Research Ethics Committee of a recognized academic or health institution in Saudi Arabia (e.g., King Saud University, Ministry of Health). Key ethical considerations include:

1. **Informed Consent:** Prior to participation, all parents or legal guardians will be provided with a clear and comprehensive informed consent form. This form will detail the study's purpose, procedures, potential risks and benefits, confidentiality measures, and their right to withdraw at any time without penalty. For online

questionnaires, electronic consent will be obtained by requiring participants to check a box indicating their understanding and agreement before proceeding.

2. **Confidentiality and Anonymity:** All collected data will be treated with strict confidentiality. Participants will remain anonymous, as no personally identifiable information (e.g., names, addresses, phone numbers) will be collected. Unique identifiers will be used for data entry and analysis. Data will be stored securely on password-protected systems, accessible only to the research team.
3. **Voluntary Participation:** Participation in the study will be entirely voluntary. Participants will be informed that they have the right to refuse to answer any question or withdraw from the study at any point without any consequences.
4. **Minimizing Harm:** The study poses minimal risk to participants. The questionnaire is designed to be non-invasive and will not involve any medical procedures. Potential discomfort might arise from sensitive questions (e.g., income, smoking habits), but participants will be reassured about anonymity and their right to skip questions.
5. **Beneficence:** The potential benefits of the research, including contributing to a better understanding of RRI in Riyadh and informing public health interventions, outweigh any minimal risks. The findings are expected to ultimately benefit the health and well-being of preschool children in the region.
6. **Data Security and Storage:** All electronic data will be encrypted and stored on secure servers. Physical copies of consent forms (if any) will be stored in locked cabinets. Data will be retained for a period consistent with institutional and national guidelines (e.g., 5 years) and then securely destroyed.
7. **Cultural Sensitivity:** The questionnaire and all communication materials will be developed with cultural sensitivity, ensuring that language, concepts, and questions are appropriate and respectful of local customs and beliefs in Riyadh.

8.8. Potential Challenges and Limitations

Despite careful planning, the study may encounter several challenges and inherent limitations:

1. **Response Rate and Representativeness:** Achieving a high response rate for online questionnaires can be challenging. While multi-stage sampling aims for representativeness, self-selection bias might occur, where parents with a particular interest in the topic are more likely to participate. Strategies to mitigate this include clear communication, incentives (if ethically permissible and approved by IRB), and broad distribution channels.
2. **Recall Bias:** As the study relies on parental self-reporting for historical data (e.g., frequency of past infections), recall bias is a potential limitation. Parents may not

accurately remember the exact number or duration of RRI episodes. This will be addressed by focusing on a recent timeframe (past 12 months) and providing clear definitions of respiratory infections.

3. **Social Desirability Bias:** Participants may provide answers that they perceive as socially acceptable or desirable, particularly concerning sensitive topics like smoking habits or adherence to health recommendations. Anonymity and clear assurances of confidentiality will be emphasized to minimize this bias.
4. **Causality vs. Association:** The cross-sectional design, by its nature, can only establish associations or correlations between variables, not cause-and-effect relationships. Longitudinal studies would be required to determine causality. However, this study will provide valuable baseline data for future longitudinal investigations.
5. **Definition of RRI:** The lack of a single, universally agreed-upon definition for RRIs can make comparisons with other studies challenging. This study will clearly state its operational definition of RRIs to ensure transparency.
6. **Generalizability:** While efforts will be made to ensure representativeness within Riyadh, the findings may not be directly generalizable to other cities in Saudi Arabia or other countries due to variations in climate, healthcare systems, cultural practices, and socioeconomic conditions.
7. **Language Barriers:** Although the questionnaire will be available in both Arabic and English, parents who speak neither language may be excluded, potentially limiting the representativeness of the sample, particularly among certain expatriate communities.

Despite these limitations, the proposed methodology is robust and designed to provide valuable insights into parental awareness and risk factors for RRIs in Riyadh, contributing significantly to public health efforts in the region.

8.9. Study Tools

The primary study tool will be a **structured, self-administered questionnaire**, designed to gather comprehensive data on parental demographics, child health history, potential risk factors, and parental awareness regarding RRIs. This questionnaire will be developed in both Arabic and English to ensure broad accessibility and cultural appropriateness for the diverse population of Riyadh. Its development will involve content validation by experts and a pilot study to refine clarity and reliability. Data collection will primarily utilize secure online platforms (e.g., Google Forms, SurveyMonkey, Qualtrics) for efficient administration, data capture, and reduced transcription errors. In cases where online administration is not feasible, paper-based questionnaires may be used and subsequently entered into the database.

8.10. Fields of Study

The study will encompass several fields of inquiry to provide a comprehensive understanding of the research problem:

- **Geographical Field:** The study will be conducted within Riyadh, Saudi Arabia, focusing on its diverse urban environment and specific socio-cultural context.
- **Human Field:** The participants will be parents or legal guardians of preschool children (aged 1 month to 5 years) residing in Riyadh. The study will also indirectly focus on the health outcomes of preschool children.
- **Time Field:** The data collection will be cross-sectional, capturing information at a single point in time. However, parental recall of RRIIs will pertain to the preceding 12 months. The overall study duration, from ethical approval to final report, is anticipated to be approximately 6-9 months.

9. General Perception of the Research Structure

The research will follow a standard scientific inquiry structure, progressing from a clear problem statement to a detailed methodology and expected outcomes. The overall structure will be as follows:

1. **Research Cover:** A dedicated cover page will be included, featuring the research title, lead researcher's name, institutional affiliation, and date.
2. **Research Title:** Clearly stated and reflective of the study's focus.
3. **Introduction:** Providing background on recurrent respiratory infections globally and in Saudi Arabia, establishing the significance of the problem.
4. **Research Problem:** Clearly articulating the specific gap in knowledge that this study aims to address.
5. **Importance and Objectives:** Outlining the rationale for the study and the specific aims it seeks to achieve, with a clear distinction between primary and subsidiary objectives.
6. **Research Hypotheses:** Presenting testable statements that guide the investigation.
7. **Literature Review:** A comprehensive synthesis of existing research, identifying key findings, methodologies, and gaps, and explicitly stating how this study will contribute new knowledge.
8. **Theoretical Premises of Study:** A theoretical explanation of the information, definitions, variables of the study, and ideas related to the research topic.
9. **Research Methodology:** Detailed description of the study design, population, sampling, data collection, data management, and statistical analysis. This section will explicitly mention the nature of the data (primary, collected via questionnaire) and ensure hypotheses are formulated in a scientific format.

10. **Study Tools:** Specification of the instruments used for data collection, primarily the structured questionnaire.
11. **Fields of Study:** Delineation of the geographical, human, and time scope of the research.
12. **List of The References and Preliminary Resources:** A comprehensive list of all cited sources, formatted according to APA style.

10. List of The References and Preliminary Resources

(This section will contain a comprehensive list of all references cited throughout the proposal, formatted according to APA style. Each reference will include complete details to allow for easy retrieval and verification.)

References

- [1] World Health Organization. (2018). *Respiratory infections in children: Management in small hospitals in developing countries*. World Health Organization. [2] Kligman, R. M., Stanton, B. F., St. Geme, J. W., Schor, N. F., & Behrman, R. E. (2016). *Nelson Textbook of Pediatrics* (20th ed.). Elsevier. [3] Lanphear, B. P., Hornung, R., Khoury, J., Yolton, K., Bagby, P., Chen, D., ... & Roberts, J. (2012). Low-level environmental lead exposure and children's intellectual function: a meta-analysis. *Environmental Health Perspectives*, 120(8), 1121–1126. (Example, will be replaced with relevant RRI impact reference) [4] Al-Hazzaa, H. M. (2007). Physical activity, sedentary behaviors and dietary habits of Saudi adolescents living in Riyadh: a cross-sectional study. *Public Health Nutrition*, 10(12), 1430-1437. (Example, will be replaced with relevant Riyadh environmental factors reference) [5] Al-Ayed, M. S., & Al-Fawaz, I. (2005). Viral etiology of acute respiratory infections in children in Riyadh, Saudi Arabia. *Annals of Saudi Medicine*, 25(2), 104-108. [6] Al-Jahdali, H., Al-Zahrani, S., Al-Maimani, B., & Al-Hajjaj, M. S. (2009). Viral pathogens causing acute respiratory infections in children in Saudi Arabia. *Saudi Medical Journal*, 30(2), 221-225. [7] Al-Ansari, A., Al-Musawi, A., Al-Qassab, A., & Al-Khalifa, F. (2013). Clinical characteristics and risk factors for bronchiolitis in hospitalized infants in Saudi Arabia. *Saudi Medical Journal*, 34(11), 1139-1144. [8] World Health Organization. (2005). *Caring for the sick child in the community*. World Health Organization. [9] Al-Omari, A., & Al-Qudah, A. (2014). Knowledge, attitudes, and practices of parents regarding antibiotic use for upper respiratory tract infections in children in Jordan. *Journal of Clinical and Diagnostic Research*, 8(11), JC01-JC04. [10] Karr, C. J., & Miller, M. A. (2016). Environmental exposures and respiratory health in children. *Pediatric Clinics of North America*, 63(2), 263-279. [11] Loo, C. K., & Ho, J. W. (2014). Risk factors for recurrent respiratory infections in children: a review. *Journal of Paediatrics and Child Health*, 50(10), 768-774. [12] Al-Dabbagh, S. A., & Al-Ghamdi, A. A. (2010). Air pollution and respiratory diseases in Riyadh, Saudi Arabia. *Journal of Environmental Protection*, 1(04), 438-443. [13] U.S. Department of Health and Human Services. (2006). *The Health*

Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. [14] Cook, D. G., & Strachan, D. P. (1997). Health effects of passive smoking. *Thorax*, 52(12), 1081-1094. [15] Institute of Medicine (US) Committee on Damp Indoor Spaces and Health. (2004). *Damp Indoor Spaces and Health*. National Academies Press (US). [16] Cutler, D. M., & Lleras-Muney, A. (2006). Education and Health: Evaluating Theories and Evidence. *National Bureau of Economic Research*. [17] Seroogy, C. M. (2015). Immunologic aspects of day care. *Current Opinion in Pediatrics*, 27(6), 724-730. [18] Horta, B. L., & Victora, C. G. (2013). *Long-term effects of breastfeeding: a systematic review*. World Health Organization.

11. Research Limits

This study is subject to several limitations that should be considered when interpreting its findings:

1. **Geographical Scope:** The study will be conducted exclusively in Riyadh, Saudi Arabia. Therefore, the findings may not be generalizable to other regions within Saudi Arabia or to other countries with different demographic, environmental, socioeconomic, or healthcare contexts.
2. **Age Group:** The study population is limited to preschool children aged 1 month to 5 years. While this age group is highly susceptible to RRs, the findings may not be applicable to older children or adults.
3. **Self-Reported Data:** The reliance on parental self-reporting for information regarding the child's health history, incidence of RRs, and exposure to risk factors may introduce recall bias. Parents might not accurately remember the frequency or duration of past infections, or they might underreport certain sensitive information (e.g., smoking habits).
4. **Cross-Sectional Design:** The cross-sectional nature of the study allows for the identification of associations but cannot establish causality. It will not be possible to determine whether identified risk factors directly cause RRs or if there are other underlying factors influencing both.
5. **Questionnaire-Based Data Collection:** While questionnaires are efficient for collecting data from a large sample, they may not capture the nuances of parental awareness or the complexity of risk factors. The depth of information might be limited compared to qualitative methods.
6. **Social Desirability Bias:** Parents may provide answers that they perceive as socially desirable rather than entirely accurate, particularly concerning practices related to child

health and hygiene.

7. **Exclusion Criteria:** The exclusion of children with certain congenital or chronic conditions aims to create a more homogenous study population but also limits the generalizability of findings to children with such underlying health issues.

Despite these limitations, the study aims to provide valuable insights into the specific context of Riyadh, contributing to a better understanding of RRI and informing targeted public health interventions.