

Probability and Statistics

Prepared by Dr. Alfaisal Hasan

STA 201
Probability and Statistics

Lecture NO 10



Probability and Statistics

Lecture NO 10

STA 201

Probability and Statistics

Prepared by
Dr. Alfaisal Hasan

Chapter

Topics

- Chapter 1: Introduction To Statistics
 - Chapter 2: Descriptive Statistics
 - Chapter 3: Probability
 - Chapter 4: Discrete Probability Distributions
 - Chapter 5: Normal Probability Distributions
 - Chapter 6: Confidence Intervals
 - Chapter 7: Hypothesis Testing with One Sample
 - Chapter 8: Hypothesis Testing with Two Samples
 - Chapter 9: Correlation And Regression
 - Chapter 10: Chi-Square Tests and The F -Distribution
- الفصل الأول: مقدمة في الإحصاء
الفصل الثاني: الإحصاء الوصفي
الفصل الثالث: الاحتمالات
الفصل الرابع: التوزيعات الاحتمالية المنفصلة
الفصل الخامس: التوزيع الاحتمالي الطبيعي
الفصل السادس: فترات الثقة
الفصل السابع: اختبار الفرضيات لعينة واحدة
الفصل الثامن: اختبار الفرضيات لعينتين
الفصل التاسع: الارتباط والانحدار
الفصل العاشر: اختبارات كاي تربيع وتوزيع F

Chapter Outline

10.1 Goodness of Fit

10.1 صلاح الملاءمة

10.2 Independence

10.2 الاستقلال

10.3 Comparing Two Variances

10.3 المقارنة بين تباينين

10.4 Analysis of Variance

10.4 تحليل التباين

Section 10.1: Goodness of Fit

A **multinomial experiment** is a probability experiment consisting of a fixed number of trials in which there are more than two possible outcomes for each independent trial. (Unlike the **binomial** experiment in which there were only two possible outcomes.)

التجربة متعددة الحدود هي تجربة احتمالية تتكون من عدد ثابت من التجارب التي يوجد فيها أكثر من نتيجتين محتملتين لكل تجربة مستقلة. (على عكس التجربة ذات الحدين التي لم يكن هناك سوى نتيجتين محتملتين).

Example:

A researcher claims that the distribution of favorite pizza toppings among teenagers is as shown below.

يدعي أحد الباحثين أن توزيع طبقة البيتزا المفضلة بين المراهقين هو كما هو موضح أدناه.

Each outcome is classified into **categories**.

Topping	Frequency, f
Cheese	41%
Pepperoni	25%
Sausage	15%
Mushrooms	10%
Onions	9%

The probability for each possible outcome is fixed.

Chi-Square Goodness of Fit test

A **Chi-Square Goodness-of-Fit Test** is used to test whether a frequency distribution fits an expected distribution.

يتم استخدام اختبار كاي تربيع لاختبار ما إذا كان التوزيع التكراري يناسب التوزيع المتوقع

To calculate the test statistic for the chi-square goodness-of-fit test, the observed frequencies and the expected frequencies are used.

لحساب إحصائية الاختبار لاختبار جودة المطابقة لمربع كاي، يتم استخدام التكرارات المرصودة والتكرارات المتوقعة.

The **observed frequency** O of a category is the frequency for the category observed in the sample data.

التردد الملحوظ لفئة ما هو تكرار الفئة الملحوظة في بيانات العينة.

The **expected frequency** E of a category is the *calculated* frequency for the category. Expected frequencies are obtained assuming the specified (or hypothesized) distribution. The expected frequency for the i th category is $E_i = np_i$

where n is the number of trials (the sample size) and p_i is the assumed probability of the i th category.

التكرار المتوقع لفئة ما هو التكرار المحسوب للفئة.

Observed and Expected Frequencies

Example:

200 teenagers are randomly selected and asked what their favorite pizza topping is. The results are shown below.

Find the observed frequencies and the expected frequencies.

يتم اختيار 200 مراهق بشكل عشوائي وسؤالهم عن الطبقة المفضلة لديهم للبيتزا. النتائج مبينة أدناه.
أوجد التكرارات المرصودة والتكرارات المتوقعة.

Topping	Results ($n = 200$)	% of teenagers
Cheese	78	41%
Pepperoni	52	25%
Sausage	30	15%
Mushrooms	25	10%
Onions	15	9%

Observed Frequency	Expected Frequency
78	$200(0.41) = 82$
52	$200(0.25) = 50$
30	$200(0.15) = 30$
25	$200(0.10) = 20$
15	$200(0.09) = 18$

Chapter 10 Chi-Square Goodness of Fit test

For the chi-square goodness-of-fit test to be used, the following must be true.

1. The observed frequencies must be obtained by using a random sample.
2. Each expected frequency must be greater than or equal to 5.

The Chi-Square Goodness-of-Fit Test

If the conditions listed above are satisfied, then the sampling distribution for the goodness-of-fit test is approximated by a chi-square distribution with $k - 1$ degrees of freedom, where k is the number of categories. The test statistic for the chi-square goodness-of-fit test is

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

The test is always a right-tailed test.

where O represents the observed frequency of each category and E represents the expected frequency of each category.

Chapter 10 Chi-Square Goodness of Fit test

Performing a Chi-Square Goodness-of-Fit Test

In Words

1. Identify the claim. State the null and alternative hypotheses.
2. Specify the level of significance.
3. Identify the degrees of freedom.
4. Determine the critical value.
5. Determine the rejection region.

In Symbols

State H_0 and H_a .

Identify α .

$$\text{d.f.} = k - 1$$

Use Table Chi Square.

Chapter 10 Chi-Square Goodness of Fit test

Performing a Chi-Square Goodness-of-Fit Test

Word	Symbol
Identify the claim. State the null and alternative hypotheses. تحديد الادعاء. اذكر الفرضيتين الصفري والبديل	State H_0 and H_a .
Specify the level of significance. تحديد مستوى المعنوية	Identify α .
Identify the degrees of freedom. تحديد درجات الحرية	d.f. = $k - 1$
Determine the critical value. تحديد القيمة الحرجة	Use Table Chi Square.
Determine the rejection region. تحديد منطقة الرفض	

Correlation coefficient

Word	Symbol
6- Calculate the test statistic. حساب القيمة الناتجة من الاختبار	$\chi^2 = \sum \frac{(O - E)^2}{E}$
Make a decision to reject or fail to reject the null hypothesis. تحديد القرار الاحصائي برفض الفرض الصفري أو الفشل في رفضه	If χ^2 is in the rejection region, reject H_0. Otherwise, fail to reject H_0.
Interpret the decision in the context of the original claim. تحديد القرار الاحصائي عن طريق الادعاء	

Chapter **Example**

A researcher claims that the distribution of favorite pizza toppings among teenagers is as shown below. 200 randomly selected teenagers are surveyed.

يدعي أحد الباحثين أن توزيع طبقة البيتزا المفضلة بين المراهقين هو كما هو موضح أدناه. تم استطلاع رأي 200 مراهق تم اختيارهم عشوائياً.

Topping	Frequency, f
Cheese	39%
Pepperoni	26%
Sausage	15%
Mushrooms	12.5%
Onions	7.5%

Using $\alpha = 0.01$, and the observed and expected values previously calculated, test the surveyor's claim using a chi-square goodness-of-fit test.

والقيم المرصودة والمتوقعة المحسوبة مسبقاً، تختبر ادعاء المساح باستخدام اختبار جودة المطابقة لمربع كاي.

Chapter 10 Solution

H_0 : The distribution of pizza toppings is 39% cheese, 26% pepperoni, 15% sausage, 12.5% mushrooms, and 7.5% onions. (Claim)

H_a : The distribution of pizza toppings differs from the claimed or expected distribution.

Because there are 5 categories, the chi-square distribution has $k - 1 = 5 - 1 = 4$ degrees of freedom.

With d.f. = 4 and $\alpha = 0.01$, the critical value is $\chi^2_0 = 13.277$.

Chapter Solution


Critical Value for χ^2_0

With d.f. = 4 and $\alpha = 0.01$

Degrees of freedom	α									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	—	—	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.145	12.838
4	0.207	0.267	0.404	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.071	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278

The critical value is $\chi^2_0 = 13.277$.

Chapter 1 Solution



Topping	Observed Frequency	Expected Frequency
Cheese	78	82
Pepperoni	52	50
Sausage	30	30
Mushrooms	25	20
Onions	15	18

$$X^2 = \sum \frac{(O - E)^2}{E} = \frac{(78 - 82)^2}{82} + \frac{(52 - 50)^2}{50} + \frac{(30 - 30)^2}{30} + \frac{(25 - 20)^2}{20} + \frac{(15 - 18)^2}{18}$$

$$\approx 2.025$$

Fail to reject H_0 .

There is not enough evidence at the 1% level to reject the surveyor's claim.

Section 10.2 Independence

An $r \times c$ **contingency table** shows the observed frequencies for two variables. The observed frequencies are arranged in r rows and c columns. The intersection of a row and a column is called a cell.

يوضح جدول الطوارئ التكرارات المرصودة لمتغيرين. يتم ترتيب الترددات المرصودة في صفوف وأعمدة يسمى تقاطع الصف والعمود بالخلية.

The following contingency table shows a random sample of 321 fatally injured passenger vehicle drivers by age and gender. (Adapted from Insurance Institute for Highway Safety)

يبين جدول الطوارئ التالي عينة عشوائية مكونة من 321 سائق سيارة ركاب مصابين بجروح قاتلة حسب العمر والجنس. (مقتبس من معهد التأمين للسلامة على الطرق السريعة)

	Age					
Gender	16 – 20	21 – 30	31 – 40	41 – 50	51 – 60	61 and older
Male	32	51	52	43	28	10
Female	13	22	33	21	10	6

Expected Frequency

Assuming the two variables are independent, you can use the contingency table to find the expected frequency for each cell.

بافتراض أن المتغيرين مستقلان، يمكنك استخدام جدول الاحتمالات للعثور على التكرار المتوقع لكل خلية.

Finding the Expected Frequency for Contingency Table Cells

The expected frequency for a cell $E_{r,c}$ in a contingency table is

إيجاد التردد المتوقع لخلايا جدول الطوارئ التكرار المتوقع للخلية في جدول الطوارئ هو

Chapter Example

Find the expected frequency for each “Male” cell in the contingency table for the sample of 321 fatally injured drivers. Assume that the variables, age and gender, are independent.

Gender	Age						Total
	16 – 20	21 – 30	31 – 40	41 – 50	51 – 60	61 and older	
Male	32	51	52	43	28	10	216
Female	13	22	33	21	10	6	105
Total	45	73	85	64	38	16	321

Chapter Example

Gender	Age						Total
	16 – 20	21 – 30	31 – 40	41 – 50	51 – 60	61 and older	
Male	32	51	52	43	28	10	216
Female	13	22	33	21	10	6	105
Total	45	73	85	64	38	16	321

Expected frequency $E_{r,c} = \frac{(\text{Sum of row } r) \times (\text{Sum of column } c)}{\text{Sample size}}$

$$E_{11} = \frac{216 \cdot 45}{321} \approx 30.28 \quad E_{12} = \frac{216 \cdot 73}{321} \approx 49.12 \quad E_{13} = \frac{216 \cdot 85}{321} \approx 57.20$$

$$E_{14} = \frac{216 \cdot 64}{321} \approx 43.07 \quad E_{15} = \frac{216 \cdot 38}{321} \approx 25.57 \quad E_{16} = \frac{216 \cdot 16}{321} \approx 10.77$$

Chi-Square Independence Test

A **chi-square independence test** is used to test the independence of two variables. Using a chi-square test, you can determine whether the occurrence of one variable affects the probability of the occurrence of the other variable.

For the chi-square independence test to be used, the following must be true.

- 1.The observed frequencies must be obtained by using a random sample.
- 2.Each expected frequency must be greater than or equal to 5.

Chi-Square Independence Test

A **chi-square independence test** is used to test the independence of two variables. Using a chi-square test, you can determine whether the occurrence of one variable affects the probability of the occurrence of the other variable.

For the chi-square independence test to be used, the following must be true.

- 1.The observed frequencies must be obtained by using a random sample.
- 2.Each expected frequency must be greater than or equal to 5.

Chapter Example

The following contingency table shows a random sample of 321 fatally injured passenger vehicle drivers by age and gender. The expected frequencies are displayed in parentheses. At $\alpha = 0.05$, can you conclude that the drivers' ages are related to gender in such accidents?

يبين جدول الطوارئ التالي عينة عشوائية مكونة من 321 سائق سيارة ركاب مصابين بجروح قاتلة حسب العمر والجنس. يتم عرض الترددات المتوقعة بين قوسين. هل يمكن استنتاج أن أعمار السائقين مرتبطة بالجنس في مثل هذه الحوادث؟

Gender	Age						Total
	16 – 20	21 – 30	31 – 40	41 – 50	51 – 60	61 and older	
Male	32 (30.28)	51 (49.12)	52 (57.20)	43 (43.07)	28 (25.57)	10 (10.77)	216
Female	13 (14.72)	22 (23.88)	33 (27.80)	21 (20.93)	10 (12.43)	6 (5.23)	105
	45	73	85	64	38	16	321

Chapter **Solution**

Because each expected frequency is at least 5 and the drivers were randomly selected, the chi-square independence test can be used to test whether the variables are independent.

H_0 : The drivers' ages are independent of gender.

H_a : The drivers' ages are dependent on gender. (Claim)

$$\text{d.f.} = (r - 1)(c - 1) = (2 - 1)(6 - 1) = (1)(5) = 5$$

With d.f. = 5 and $\alpha = 0.05$, the critical value is $\chi^2_0 = 11.071$.

Chapter Solution

Critical Value for χ^2_0


$$\text{d.f.} = (r-1)(c-1) = (2-1)(6-1) = (1)(5) = 5 \quad \text{and} \quad \alpha = 0.05$$

Degrees of freedom	α									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	—	—	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.879	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.601	9.236	11.071	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188

The critical value is $\chi^2_0 = 11.071$

Chapter 10 Solution

Example continued:



$$X^2 = \sum \frac{(O - E)^2}{E} = 2.84$$

Fail to reject H_0 .

O	E	$O - E$	$(O - E)^2$	$\frac{(O - E)^2}{E}$
32	30.28	1.72	2.9584	0.0977
51	49.12	1.88	3.5344	0.072
52	57.20	-5.2	27.04	0.4727
43	43.07	-0.07	0.0049	0.0001
28	25.57	2.43	5.9049	0.2309
10	10.77	-0.77	0.5929	0.0551
13	14.72	-1.72	2.9584	0.201
22	23.88	-1.88	3.5344	0.148
33	27.80	5.2	27.04	0.9727
21	20.93	0.07	0.0049	0.0002
10	12.43	-2.43	5.9049	0.4751
6	5.23	0.77	0.5929	0.1134

There is not enough evidence at the 5% level to conclude that age is dependent on gender in such accidents.

10.3 Comparing Two Variances

Let s_1^2 and s_2^2 represent the sample variances of two different populations. If both populations are normal and the population variances σ_1^2 and σ_2^2 are equal, then the sampling distribution of

$$F = \frac{s_1^2}{s_2^2}$$

is called an ***F*-distribution**.


There are several properties of this distribution.

1. The *F*-distribution is a family of curves each of which is determined by two types of degrees of freedom: the degrees of freedom corresponding to the variance in the numerator, denoted **d.f._N**, and the degrees of freedom corresponding to the variance in the denominator, denoted **d.f._D**. Continued.

F-Distribution

Properties of the F -distribution continued:

2. F -distributions are positively skewed.
3. The total area under each curve of an F -distribution is equal to 1.
4. F -values are always greater than or equal to 0.
5. For all F -distributions, the mean value of F is approximately equal to 1.



Critical Values for the F-Distribution

Finding Critical Values for the F -Distribution

1. Specify the level of significance α .
2. Determine the degrees of freedom for the numerator, d.f._N.
3. Determine the degrees of freedom for the denominator, d.f._D.
4. Use Table 7 in Appendix B to find the critical value. If the hypothesis test is
 - a. one-tailed, use the α F -table.
 - b. two-tailed, use the $\alpha/2$ F -table.

Chapter Example

Find the critical F -value for a right-tailed test when $\alpha = 0.05$, $d.f._N = 5$ and $d.f._D = 28$.

F -Distribution Table

d.f. _D : Degrees of freedom, denominator	$\alpha = 0.05$					
	d.f. _N : Degrees of freedom, numerator					
	1	2	3	4	5	6
1	161.4	199.5	215.7	224.6	230.2	234.0
2	18.51	19.00	19.16	19.25	19.30	19.33
//						
27	4.21	3.35	2.96	2.73	2.57	2.46
→ 28	4.20	3.34	2.95	2.71	2.56	2.45
29	4.18	3.33	2.93	2.70	2.55	2.43

The critical value is $F_0 = 2.56$.

Chapter Example

Find the critical F -value for a two-tailed test when $\alpha = 0.10$, $d.f._N = 4$ and $d.f._D = 6$.

$$\alpha = (0.10) = 0.05$$

Appendix B: Table 7: F -Distribution

d.f. _D : Degrees of freedom, denominator	$\alpha = 0.05$					
	d.f. _N : Degrees of freedom, numerator					
	1	2	3	4	5	6
1	161.4	199.5	215.7	224.6	230.2	234.0
2	18.51	19.00	19.16	19.25	19.30	19.33
3	10.13	9.55	9.28	9.12	9.01	8.94
4	7.71	6.94	6.59	6.39	6.26	6.16
5	6.61	5.79	5.41	5.19	5.05	4.95
→ 6	5.99	5.14	4.76	4.53	4.39	4.28
7	5.59	4.74	4.35	4.12	3.97	3.87

The critical value is $F_0 = 4.53$.

Two-Sample F-Test for Variances

Two-Sample F -Test for Variances

A two-sample F-test is used to compare two population variances σ_1^2 and σ_2^2 when a sample is randomly selected from each population. The populations must be independent and normally distributed. The test statistic is

$$F = \frac{s_1^2}{s_2^2}$$

where s_1^2 and s_2^2 represent the sample variances with $s_1^2 \geq s_2^2$. The degrees of freedom for the numerator is d.f._N = $n_1 - 1$ and the degrees of freedom for the denominator is d.f._D = $n_2 - 1$, where n_1 is the size of the sample having variance s_1^2 and n_2 is the size of the sample having variance s_2^2 .

Chapter **Example**

A travel agency's marketing brochure indicates that the standard deviations of hotel room rates for two cities are the same. A random sample of 13 hotel room rates in one city has a standard deviation of \$27.50 and a random sample of 16 hotel room rates in the other city has a standard deviation of \$29.75. Can you reject the agency's claim at $\alpha = 0.01$?


Because $29.75 > 27.50$, $s_1^2 = 885.06$ and $s_2^2 = 756.25$.

$$H_0: \sigma_1^2 = \sigma_2^2 \quad (\text{Claim})$$

$$H_a: \sigma_1^2 \neq \sigma_2^2$$

Solution

This is a two-tailed test with $\alpha = 0.01 = 0.005$,
 $d.f._N = 15$ and $d.f._D = 12$.



The critical value is $F_0 = 4.72$.

The test statistic is

$$F = \frac{s_1^2}{s_2^2} = \frac{885.06}{756.25} \approx 1.17.$$

Fail to reject H_0 .

There is not enough evidence at the 1% level to reject the claim that the standard deviation of the hotel room rates for the two cities are the same.

10.4 Analysis of Variance (ANOVA)

One-way analysis of variance is a hypothesis-testing technique that is used to compare means from three or more populations. Analysis of variance is usually abbreviated **ANOVA**.

تحليل التباين أحادي الاتجاه هو أسلوب لاختبار الفرضيات يستخدم لمقارنة المتوسطات من ثلاثة مجتمعات أو أكثر.
عادة ما يتم اختصار تحليل التباين

In a one-way ANOVA test, the following must be true:

1. Each sample must be randomly selected from a normal, or approximately normal, population.
2. The samples must be independent of each other.
3. Each population must have the same variance.

Analysis of Variance (ANOVA)

$$\text{Test statistic} = \frac{\text{Variance between samples}}{\text{Variance within samples}}$$

1. The variance between samples MS_B measures the differences related to the treatment given to each sample and is sometimes called the **mean square between**.

1. The variance within samples MS_W measures the differences related to entries within the same sample. This variance, sometimes called the **mean square within**, is usually due to sampling error.

Analysis of Variance (ANOVA)

One-Way Analysis of Variance Test

If the conditions listed are satisfied, then the sampling distribution for the test is approximated by the F-distribution. The test statistic is

$$F = \frac{MS_B}{MS_W}.$$

The degrees of freedom for the F -test are

$$\text{d.f.}_N = k - 1$$

and

$$\text{d.f.}_D = N - k$$

where k is the number of samples and N is the sum of the sample sizes.

ANOVA summary Table

A table is a convenient way to summarize the results in a one-way ANOVA test.

Variation	Sum of squares	Degrees of freedom	Mean squares	F
Between	SS_B	d.f. _N	$MS_B = \frac{SS_B}{\text{d.f.}_N}$	$MS_B \div MS_W$
Within	SS_W	d.f. _D	$MS_W = \frac{SS_W}{\text{d.f.}_D}$	

Chapter Example

The following table shows the salaries of randomly selected individuals from four large metropolitan areas. At $\alpha = 0.05$, can you conclude that the mean salary is different in at least one of the areas? (Adapted from US Bureau of Economic Analysis)

Pittsburgh	Dallas	Chicago	Minneapolis
27,800	30,000	32,000	30,000
28,000	33,900	35,800	40,000
25,500	29,750	28,000	35,000
29,150	25,000	38,900	33,000
30,295	34,055	27,245	29,805

1- State H_0 and H_a

2- Critical value is F_0

3- Test statistics F using ANOVA table

Chapter **Solution**

$$H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$$

H_a : At least one mean is different from the others. (Claim)

Because there are $k = 4$ samples, $\text{d.f.}_N = k - 1 = 4 - 1 = 3$.

The sum of the sample sizes is

$$N = n_1 + n_2 + n_3 + n_4 = 5 + 5 + 5 + 5 = 20.$$

$$\text{d.f.}_D = N - k = 20 - 4 = 16$$

Using $\alpha = 0.05$, $\text{d.f.}_N = 3$, and $\text{d.f.}_D = 16$,
the critical value is $F_0 = 3.24$.

Chapter Solution

To find the test statistic, the following must be calculated.

$$\bar{\bar{x}} = \frac{\sum x}{N} = \frac{140745 + 152705 + 161945 + 167805}{20} = 31160$$

Pittsburgh	Dallas	Chicago	Minneapolis
27,800	30,000	32,000	30,000
28,000	33,900	35,800	40,000
25,500	29,750	28,000	35,000
29,150	25,000	38,900	33,000
30,295	34,055	27,245	29,805

$$SS_B = \sum n_i(\bar{x} - \bar{\bar{x}})^2 =$$

$$= 5(28149 - 31160)^2 + 5(30541 - 31160)^2 + 5(32389 - 31160)^2 + 5(33561 - 31160)^2$$

$$SS_B = 83622620.01$$

$$MS_B = \frac{SS_B}{d.f.N} = \frac{83622.620.01}{k-1} = \frac{83622.620.01}{4-1} = 27874206.67$$

Chapter Solution

$$SS_W = \sum (n_i - 1) s_i^2$$

$$= (5 - 1)(3192128.94) + (5 - 1)(13813030.08) + (5 - 1)(24975855.83) + (5 - 1)(17658605.02)$$

$$= 238558479.52$$

$$MS_W = \frac{SS_W}{d.f.D} = \frac{238558479.52}{N-k} = \frac{238558479.52}{20-4} = 14909904.97$$

Chapter **Solution**

$$F = \frac{MS_B}{MS_W} = \frac{27874206.67}{14909904.34} \approx 1.870$$

Test
statisticCritical
value

$$1.870 < 3.24.$$

Fail to reject H_0 .

There is not enough evidence at the 5% level to conclude that the mean salary is different in at least one of the areas.

Chapter Solution

Step 1:

$$H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$$

H_a : At least one mean is different from the others. (Claim)

Step 2: The Critical value $F_0=3.24$ (From F Table)

Step 3: Test statistic $F = \frac{MS_B}{MS_W} = 1.870$

Step 4: Decision

Test
statistic Critical
value

$$1.870 < 3.24.$$


Fail to reject H_0 .

Chapter 10 Section 10.1: Goodness of Fit

Variation	Sum of Squares	Degrees of freedom	Mean Squares	F
Between				
Within				

AI in Mathematics

Chapter




الأسس
الرياضية
للحاسب


AI in Mathematics

Chapter


الخوارزميات



Algorithm



AI in Mathematics



علم البيانات

Data Analysis تحليل البيانات

علم التشفير




**On some Algebraic
methods in Cryptography**

يستخدم التشفير
مفاهيم رياضية
معقدة لضمان أمان
المعلومات
وحمايتها من
الوصول غير المصرح
به. تشمل
التطبيقات عملية
تشفير البيانات وفك
تشفيرها بواسطة
خوارزميات رياضية
معترف بها.

AI in Mathematics

المحاكاة

تعتبر الرياضيات التطبيقية
ضرورية لتطوير نماذج
رياضية تحاكي الواقع
وتساعد في حل المشكلات
العملية. تشمل تطبيقاتها
في الحوسبة تطوير
البرمجيات والنمذجة
الرياضية.



AI in Mathematics

الذكاء الاصطناعي
هو قدرة الأنظمة
البرمجية على محاكاة
وظائف التفكير
البشري مثل التعلم،
الاستدلال، واتخاذ
القرارات. يتم
استخدامه في
مجالات علمية
وحياتية متعددة

AI in Mathematics



AI in Mathematics

تخصيص التعليم

يمكن للذكاء الاصطناعي
تخصيص مناهج الرياضيات
لتناسب مستوى كل طالب،
مما يساعد على تعزيز الفهم
واستيعاب المفاهيم.
التطبيقات الذكية تستخدم
لتحديد نقاط القوة والضعف
لدى الطلاب وتقديم محتوى
مخصص.



Chapter Summary

Chapter

10.1 Goodness of Fit (**Chi-Square**)

10.2 Independence (**Chi-Square**)

10.3 Comparing Two Variances (**F-distribution**)

10.4 Analysis of Variance (**F-distribution**)

Un Solved Examples

Chapter

In Chi-Square test, If the critical value $\chi_0^2 = 14.067$ and the test statistic $X^2 = 7.13$, then we reject H_0

في اختبار كاي سكوير إذا كانت القيمة الحرجة $\chi_0^2 = 14.067$ والقيمة الناتجة من الاختبار $X^2 = 7.13$ فإننا نرفض الفرض الصفري

Un Solved Examples

Chapter

In Anova test, If the critical value = 3.52 and test statistic $F = 1.81$, then we reject H_0

في اختبار التباين أنوفا إذا كانت القيمة الحرجة = 3.52 و القيمة الناتجة من الاختبار $F = 1.81$ فإننا نرفض الفرض الصفري



Thank You