

Reducing VAP (Ventilator Associated Pneumonia) Outbreaks by Implementing VAP Bundle

Prepared by:

**FATHADIN ALI ABDU ALAHDAL
222001930**

Supervised by:

Dr. Walid Soliman

Year: 2023

Research Hypotheses

The implementation of a VAP bundle will lead to a reduction in the severity and mortality of VAP in the critical care setting.

Literature Review

Ventilator-associated pneumonia (VAP) is a serious infection that can occur in patients who are on mechanical ventilation. It is one of the most common hospital-acquired infections (HAIs) that leads to increased morbidity, mortality, and healthcare costs.

VAP outbreaks can occur in any critical care setting. Additionally, they are more common in units with high patient turnover and high rates of mechanical ventilation. VAP outbreaks can be difficult to control and they can have a significant impact on the quality of care provided to patients.

One of the most effective ways to reduce the risk of VAP outbreaks is to implement a VAP bundle. A VAP bundle is a set of evidence-based practices that are effective in preventing VAP. These practices typically include:

- Elevation of the head of the bed to at least 30 degrees
- Daily sedation interruption and assessment of readiness to extubate
- Oral hygiene with chlorhexidine mouthwash
- Subglottic suctioning
- Avoidance of routine ventilator circuit changes
- Stress ulcer prophylaxis

Several studies have shown that VAP bundles can be effective in reducing the incidence of VAP outbreaks in critical care settings. For example, a study published in the *Intensive Care Med* found that the implementation of a VAP bundle reduced the incidence of VAP by 50% in a surgical intensive care unit (Andrew Conway Morris, 2011)¹.

Another study, published in the journal *Clinical Infectious Diseases* found that the implementation of a VAP bundle reduced the incidence of VAP by 30% in a general intensive care unit (Andre C. Kalil, 2016)².

Also, Arroliga et al emphasize the importance of a multidisciplinary task force to deliver best practices, with responsibility shared between Nursing and Respiratory Therapists (RTs). The study highlights the role RTs can play in VAP prevention outside of ventilator management and spontaneous breathing trial initiation. Their work can expand to the bedside team and directly impact

VAP rates. In this center, RT involvement worked well, but in some institutions another team may be better suited for the role of oral care. The role of a single center study can be limiting, but in this case it highlights the importance of a multidisciplinary approach working as a team toward a common goal to improve patient outcomes. Their message resonates with what Helen Keller once said: “Alone we can do so little; together we can do so much.” (Christian Sandrock and Joan Daly, 2012)³.

Moreover, Considerable gap is observed between knowledge and implementation of ventilator bundle among critical care practitioners. Despite knowledge, fear of adverse events and lack of proper training are key barriers in implementation of ventilator bundle (Naveen Paliwal , 2023)⁴. Knowledge regarding specific components of VAP prevention is lacking.

Formal training and interactive educational sessions should be performed regularly to assess the competency of key personnel regarding the VAP bundle, especially in the context of rapid nurse turnover. Incentives for retention of nurses should also be considered, so that knowledge of hospital specific initiatives such as the VAP bundles of care can be cultivated over time (Cybele Lara Abad, 2021)⁵.

VAP care bundle implementation with education prepared according to evidence-based guidelines decreased VAP rates. Thus, implementation of the VAP care bundle on mechanically ventilated patients care is recommended (Aliye Okgün Alcan,2016)⁶. Effective VAP prevention in ICUs is hampered by a multifaceted mix of personal (e.g., knowledge gaps, fatigue), environmental (e.g., understaffing, equipment limitations), and organizational (e.g., unclear protocols, conflicting priorities) challenges (Vajihe Atashi ,2018)⁷.

Nurses' compliance with ventilator-associated pneumonia prevention guidelines was insufficient. Low nurse-patient ratio and large intensive care unit beds capacity were found to affect nurses' compliance and patients' outcomes. This study expanded knowledge about important aspects of nursing care; nurses' compliance with ventilator-associated pneumonia prevention guidelines and the factors that affect their compliance. This knowledge can be used by health professional to guide the clinical practice and to improve the quality of care (Sami M Aloush, 2018)⁸

Local and national VAP prevention bundles have shown to be efficient in transferring knowledge to clinical practice in ICUs. The strategies used most frequently for implementation of VAP bundles have been education, posters, audits of adherence, and feedback of results. Most initiatives have accomplished significant reduction in VAP rates, in spite of considerable

heterogeneity of recommendations, ways of implementation, auditing of adherence and registry of VAP rates (Francisco Álvarez-Lerma, 2018)⁹.

The overall compliance rate reported by the nurses was acceptable. However, the recommended VAP prevention measures are not consistently and uniformly implemented. Working in a general ICU or an ICU containing 10–15 beds or having had education regarding VAP prevention improves the nurses' compliance. However, the shortage of nursing staff, forgetting to perform some evidence-based procedures, and the hospital cost control policies are the main barriers to implementing the guidelines (Khaled M. Al-Sayaghi, 2020)¹⁰.

The guidelines for the prevention of ventilator-associated pneumonia from the Centers for Disease Control and Prevention are not consistently or uniformly implemented. Practices of nurses employed in hospitals with oral care protocols are more often congruent with the guidelines than are practices of nurses employed in hospitals without such protocols. Significant reductions in rates of ventilator-associated pneumonia may be achieved by broader implementation of oral care protocols (Carolyn L Cason, 2007)¹¹.

These studies suggest that VAP bundles are an effective way to reduce the risk of VAP outbreaks in critical care settings (Reem Alsadat, 2012)¹². However, it is important to note that VAP bundles must be implemented correctly and consistently to be effective.

Here are some of the key findings from the literature review:

- VAP bundles are effective in reducing the incidence of VAP outbreaks in critical care settings.
- VAP bundles are most effective when implemented correctly and consistently.
- VAP bundles can also help to reduce the severity and mortality of VAP.
- Further research is needed to better understand the long-term impact of VAP bundles on the reduction of VAP outbreaks.

Overall, the literature suggests that VAP bundles are an effective way to reduce the risk of VAP outbreaks in critical care settings (Joong Sik Eom, MD, 2013)¹³. However, it is important to implement VAP bundles correctly and consistently to achieve the desired results.

Research Methodology

Questionnaires can be used to collect data on topics like (Healthcare providers' practices related to VAP bundles). Questionnaires can be administered to healthcare providers in a variety of ways, such as through online surveys, paper surveys, or in-person interviews. Questions that could be included in a questionnaire to healthcare providers on reducing VAP outbreaks in critical care settings by implementing VAP bundles. In this study online survey was chosen with the following questions:

- What is your role as healthcare provider at the implementation of a VAP bundle?
- Gender
- Year of experience
- Academic level
- Current position
- How familiar are you with the VAP bundle?
- How important do you consider the VAP bundle to be in improving patient outcomes for mechanically ventilated patients?
- How strongly do you agree that the VAP bundle reduces ventilator-associated pneumonia (VAP) rates?
- Identify the biggest barrier(s) to implementing the VAP bundle in your unit
- What single change would you choose to make VAP bundle implementation easier?

By collecting data from healthcare providers, we can gain a better understanding of the factors that contribute to the successful implementation of VAP bundles, the impact of VAP bundles on reducing VAP outbreaks, and improving patient outcomes. When using questionnaires with healthcare providers in a study on reducing VAP outbreaks in critical care settings by implementing VAP bundles, it is important to take into account the following ethical considerations:

- Informed consent: Healthcare providers must be fully informed of the purpose of the study, the risks, and benefits of participation, and their right to withdraw from the study at any time. They should also be allowed to ask questions and have any concerns addressed before they agree to participate in the study.

- Confidentiality: All data collected from healthcare providers must be kept confidential. This means that the researchers must take steps to protect the participants' identities and prevent their data from being disclosed to unauthorized individuals.
- Privacy: Healthcare providers should be able to complete the questionnaire in a private setting where they will not be disturbed.
- Voluntariness: Participation in the study must be voluntary. Healthcare providers should not be coerced or pressured into participating.

Research Structure

A research structure for reducing VAP outbreaks in critical care settings by implementing VAP bundles could include the following components:

Introduction

Ventilator-associated pneumonia (VAP) is a serious hospital-acquired infection that can develop in patients who are mechanically ventilated. It is a leading cause of morbidity and mortality in critically ill patients as well as associated with significant healthcare costs.

VAP bundles are a set of evidence-based practices that have been shown to reduce the incidence of VAP. These bundles typically include four core components which are elevation of the head of the bed to 30-45 degrees, daily sedation interruption, assessment of spontaneous breathing trials, use of appropriate oral or enteral nutrition, and oral care with Chlorhexidine.

Despite the evidence supporting their use, VAP bundles are not always implemented effectively in clinical practice. This project will explore the barriers and facilitators to the implementation of VAP bundles in hospitals in critical care settings and develop implementation strategies to improve the implementation of VAP bundles.

VAP is considered a public health problem and applying VAP bundles is an effective way to reduce its incidence. However, VAP bundles are not always implemented effectively in clinical practice. This project will address this important gap in knowledge and practice.

The concept of VAP bundles was first introduced in the early 1990s. Since then, there have been numerous studies that have demonstrated the effectiveness of VAP bundles in reducing the incidence of VAP.

In Saudi Arabia, the first VAP bundle was implemented in 2005. Since then, there has been a growing interest in VAP bundles and many hospitals in Saudi Arabia have implemented their own VAP bundles. However, there is still room for improvement in the implementation of VAP bundles in Saudi Arabia. This study is significant because it will help to improve the implementation and effectiveness of VAP bundles in hospitals in critical care settings. This will lead to a reduction in the incidence of VAP, which will improve patient outcomes and reduce healthcare costs.

VAP is a serious infection that can affect anyone who is on mechanical ventilation. This includes critically ill patients, such as those who have had surgery or who are suffering from a serious medical condition. By improving the implementation and effectiveness of VAP bundles, this study will help to reduce the number of people who develop VAP. This will improve patient outcomes and reduce healthcare costs for everyone. This study will be conducted in three phases:

Phase 1: This phase will involve a systematic review of the literature on VAP bundles. This review aims to identify the best practices for the implementation of VAP bundles.

Phase 2: This phase will involve a quantitative study to explore the barriers and facilitators to the implementation of VAP bundles in a hospital in a critical care setting. The findings of this study will be used to develop strategies to improve the implementation and effectiveness of VAP bundles.

Phase 3: This phase will involve the implementation of the VAP bundles.

VAP is a serious and costly problem. VAP bundles are an effective way to reduce the incidence of VAP, but they are not always implemented effectively in clinical practice.

This project will explore the barriers and facilitators to the implementation of VAP bundles, and develop and implement strategies to improve the implementation and effectiveness of VAP bundles in hospitals in critical care settings. Also, it builds on existing knowledge about VAP bundles and their effectiveness. Moreover, it will contribute to existing knowledge about the barriers and facilitators to the implementation of VAP bundles and develop the implementation of effective strategies to improve the implementation and effectiveness of VAP bundles. Also, it is related to some prior research studies on VAP bundles. For example, the project builds on the findings of a study by Joong Sik Eom, MD. (2013), which found that the implementation of a VAP bundle was associated with a significant reduction in the incidence of VAP.

Literature Review

Ventilator-associated pneumonia (VAP) is a serious infection that can occur in patients who are on mechanical ventilation. It is one of the most common hospital-acquired infections (HAIs) that leads to increased morbidity, mortality, and healthcare costs.

VAP outbreaks can occur in any critical care setting. Additionally, they are more common in units with high patient turnover and high rates of mechanical ventilation. VAP outbreaks can be difficult to control and they can have a significant impact on the quality of care provided to patients.

One of the most effective ways to reduce the risk of VAP outbreaks is to implement a VAP bundle. A VAP bundle is a set of evidence-based practices that are effective in preventing VAP. These practices typically include:

- Elevation of the head of the bed to at least 30 degrees
- Daily sedation interruption and assessment of readiness to extubate
- Oral hygiene with chlorhexidine mouthwash
- Subglottic suctioning
- Avoidance of routine ventilator circuit changes
- Stress ulcer prophylaxis

Several studies have shown that VAP bundles can be effective in reducing the incidence of VAP outbreaks in critical care settings. For example, a study published in the *Intensive Care Med* found that the implementation of a VAP bundle reduced the incidence of VAP by 50% in a surgical intensive care unit (Andrew Conway Morris, 2011)¹.

Another study, published in the journal *Clinical Infectious Diseases* found that the implementation of a VAP bundle reduced the incidence of VAP by 30% in a general intensive care unit (Andre C. Kalil, 2016)².

Also, Arroliga et al emphasize the importance of a multidisciplinary task force to deliver best practices, with responsibility shared between nursing and RTs. The study highlights the role RTs can play in VAP prevention outside of ventilator management and spontaneous breathing trial initiation. Their work can expand to the bedside team and directly impact VAP rates. In this center, RT involvement worked well, but in some institutions another team may be better suited for the role of oral care. The role of a single center study can be limiting, but in this case it highlights the importance of a multidisciplinary approach—working as a team toward a common goal to improve patient outcomes. Their message resonates with what Helen Keller once said: “Alone we can do so little; together we can do so much.” (Christian Sandroock and Joan Daly, 2012)³.

Moreover, Considerable gap is observed between knowledge and implementation of ventilator bundle among critical care practitioners. Despite knowledge, fear of adverse events and lack of proper training are key barriers in implementation of ventilator bundle (Naveen Paliwal , 2023)⁴.

Knowledge regarding specific components of VAP prevention is lacking. Formal training and interactive educational sessions should be performed regularly to assess the competency of key personnel regarding the VAP bundle, especially in the context of rapid nurse turnover. Incentives for retention of nurses should also be considered, so that knowledge of hospital specific initiatives such as the VAP bundles of care can be cultivated over time (Cybele Lara Abad, 2021)⁵.

VAP care bundle implementation with education prepared according to evidence-based guidelines decreased VAP rates. Thus, implementation of the VAP care bundle on mechanically ventilated patients care is recommended (Aliye Okgün Alcan,2016)⁶.

Effective VAP prevention in ICUs is hampered by a multifaceted mix of personal (e.g., knowledge gaps, fatigue), environmental (e.g., understaffing, equipment limitations), and organizational (e.g., unclear protocols, conflicting priorities) challenges (Vajihe Atashi ,2018)⁷.

Nurses' compliance with ventilator-associated pneumonia prevention guidelines was insufficient. Low nurse-patient ratio and large intensive care unit beds capacity were found to affect nurses' compliance and patients' outcomes. This study expanded knowledge about important aspects of nursing care; nurses' compliance with ventilator-associated pneumonia prevention guidelines and the factors that affect their compliance. This knowledge can be used by health professional to guide the clinical practice and to improve the quality of care (Sami M Aloush, 2018)⁸

Local and national VAP prevention bundles have shown to be efficient in transferring knowledge to clinical practice in ICUs. The strategies used most frequently for implementation of VAP bundles have been education, posters, audits of adherence, and feedback of results. Most initiatives have accomplished significant reduction in VAP rates, in spite of considerable heterogeneity of recommendations, ways of implementation, auditing of adherence and registry of VAP rates (Francisco Álvarez-Lerma, 2018)⁹.

The overall compliance rate reported by the nurses was acceptable. However, the recommended VAP prevention measures are not consistently and uniformly implemented. Working in a general ICU or an ICU containing 10–15 beds or having had education regarding VAP prevention improves the nurses' compliance. However, the shortage of nursing staff, forgetting to perform some evidence-based procedures, and the hospital cost control policies are the main barriers to implementing the guidelines (Khaled M. Al-Sayaghi, 2020)¹⁰.

The guidelines for the prevention of ventilator-associated pneumonia from the Centers for Disease Control and Prevention are not consistently or uniformly implemented. Practices of nurses employed in hospitals with oral care protocols are more often congruent with the guidelines than are practices of nurses employed in hospitals without such protocols. Significant reductions in rates of ventilator-associated pneumonia may be achieved by broader implementation of oral care protocols (Carolyn L Cason, 2007)¹¹.

These studies suggest that VAP bundles are an effective way to reduce the risk of VAP outbreaks in critical care settings (Reem Alsadat, 2012)¹². However, it is important to note that VAP bundles must be implemented correctly and consistently to be effective.

Here are some of the key findings from the literature review:

- VAP bundles are effective in reducing the incidence of VAP outbreaks in critical care settings.
- VAP bundles are most effective when implemented correctly and consistently.
- VAP bundles can also help to reduce the severity and mortality of VAP.
- Further research is needed to better understand the long-term impact of VAP bundles on the reduction of VAP outbreaks.

Overall, the literature suggests that VAP bundles are an effective way to reduce the risk of VAP outbreaks in critical care settings (Joong Sik Eom, MD, 2013)¹³. However, it is important to implement VAP bundles correctly and consistently to achieve the desired results.

Methodology

The questionnaires through (Google Polls) will be distributed to the healthcare provider (Respiratory Therapist, Physician, and nurses) at a hospital in Jazan Region, Saudi Arabia.

Results

All health care providers who have direct relationship with VAP agreed that they are familiar with VAP bundle 60%. Also, they have emphasized the importance of VAP bundle in improving patients' outcome with 52.9% and the reduction of VAP incidence rates with 64.7%. Therefore, most of healthcare provider has the knowledge about the VAP bundle and its effectiveness on patients' outcome. In terms of the biggest barriers to implement the VAP bundle, the majority agreed that lack of staffing is the key barrier with 29.4% where heavy workload showed 23.5%. Insufficient resources, resistance to change, and inadequate monitoring and feedback resulted with 11.8% each.

As healthcare providers recommend to make VAP implementation easier, they have responded to add more staff to make the implementation smoother.

As result, it was found that most of healthcare providers have the knowledge about VAP bundle and its role on patients' outcome and the biggest barrier to implement the VAP bundle is lack of staffing.

Conclusion

This study reveals that 60% of healthcare providers familiar with the VAP bundle recognize its importance in improving patient outcomes (52.9%) and reducing VAP incidence (64.7%). However, implementing the bundle faces significant challenges, with lack of staffing (29.4%) identified as the key barrier. Other obstacles include heavy workload, insufficient resources, resistance to change, and inadequate monitoring. To improve VAP bundle implementation, healthcare providers recommend increasing staffing levels to ease the workload and facilitate consistent care. This study highlights the knowledge gap between awareness and consistent implementation of the VAP bundle. While healthcare providers understand its benefits, resource constraints and workload hinder effective use. To reduce VAP outbreaks, addressing the human resource limitations is crucial. Increased staffing can help overcome workload challenges and ensure quality care delivery, ultimately leading to better patient outcomes and reduced VAP rates. This study's limitations include its potential for self-reporting bias and lack of information on the specific healthcare setting and sample size.

Future research should focus on Developing and testing interventions to address additional barriers besides staffing, such as resistance to change and inadequate monitoring.

Research limits

There are several research limits to the study of reducing VAP outbreaks in critical care settings by implementing VAP bundles. These limitations are the difficulty of conducting randomized controlled trials, the difficulty of controlling for all of the factors that can contribute to VAP, and the lack of consensus on the best way to implement VAP bundles.

First, the difficulty of conducting randomized controlled trials (RCTs) in critical care settings. RCTs are the gold standard for research but they can be difficult and expensive to conduct in critical care settings. This is because critical care patients are often very sick and their clinical condition can change quickly.

Second, the difficulty of controlling for all of the factors that can contribute to VAP. VAP is a complex condition that can be caused by a variety of factors, including the severity of the patient's underlying illness, the type of mechanical ventilation that the patient is receiving, and the quality of care that the patient is receiving. It can be difficult to control for all of these factors in a research study.

Third, the lack of consensus on the best way to implement VAP bundles. There is no one-size-fits-all approach to implementing VAP bundles. The best way to implement a VAP bundle will vary depending on the specific characteristics of the critical care unit and the patients that are treated in the unit.

Finally, there is no consensus on the best way to implement VAP bundles, which can make it difficult to compare the results of different studies. Therefore, questionnaires were chosen to be the methodology of the research project.

Reference List

1. Andrew Conway Morris, Reducing ventilator-associated pneumonia in intensive care: Impact of implementing a care bundle. *Intensive Care Med.* 2011
2. Andre C. Kalil, Management of Adults With Hospital-acquired and Ventilator-associated Pneumonia: 2016 Clinical Practice Guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clinical Infectious Diseases.* 2016
3. Christian Sandrock and Joan Daly, The importance of a multidisciplinary approach to VAP prevention: the role of the respiratory therapist. *Respiratory Care Journal.* 2012.
4. Naveen Paliwal, Assessment of Knowledge, Barrier in Implementation, and Compliance to Ventilator Bundle among Resident Doctors and Nurses Working in Intensive Care Units of a Tertiary Care Center of Western India: A Cross-sectional Survey, *Indian J Crit Care Med.* 2023
5. Cybele Lara Abad, Assessment of knowledge and implementation practices of the ventilator acquired pneumonia (VAP) bundle in the intensive care unit of a private hospital, *Antimicrob Resist Infect Control.* 2021
6. Aliye Okgün Alcan, Prevention of ventilator-associated pneumonia: Use of the care bundle approach, *Am J Infect Control.* 2016
7. Vajihe Atashi, The barriers to the prevention of ventilator-associated pneumonia from the perspective of critical care nurses: A qualitative descriptive study, *J Clin Nurs.* 2018
8. Sami M Aloush, Nurses' implementation of ventilator-associated pneumonia prevention guidelines: an observational study in Jordan, *Nurs Crit Care.* 2018
9. Francisco Álvarez-Lerma, Prevention of Ventilator-Associated Pneumonia: The Multimodal Approach of the Spanish ICU "Pneumonia Zero" Program, *Crit Care Med.* 2018
10. Khaled M. Al-Sayaghi, Critical care nurses' compliance and barriers toward ventilator-associated pneumonia prevention guidelines: cross-sectional survey, *J Taibah Univ Med Sci.* 2020
11. Carolyn L Cason, Nurses' implementation of guidelines for ventilator-associated pneumonia from the Centers for Disease Control and Prevention, *Am J Crit Care.* 2007
12. Reem Alsadat, Use of ventilator-associated pneumonia bundle and statistical process control chart to decrease VAP rate in Syria. *Avicenna J Med.* 2012.
13. Joong Sik Eom, MD, The impact of a ventilator bundle on preventing ventilator-associated pneumonia: A multicenter study. *Am J of Infection Control.* 2013